

9/670096

Welcome to STN International! Enter x:x

LOGINID: ssspta1813nxxm

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?): 2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Apr 08 "Ask CAS" for self-help around the clock  
NEWS 3 Jun 03 New e-mail delivery for search results now available  
NEWS 4 Aug 08 PHARMAMarketLetter(PHARMAML) - new on STN  
NEWS 5 Aug 19 Aquatic Toxicity Information Retrieval (AQUIRE)  
now available on STN  
NEWS 6 Aug 26 Sequence searching in REGISTRY enhanced  
NEWS 7 Sep 03 JAPIO has been reloaded and enhanced  
NEWS 8 Sep 16 Experimental properties added to the REGISTRY file  
NEWS 9 Sep 16 CA Section Thesaurus available in CAPLUS and CA  
NEWS 10 Oct 01 CASREACT Enriched with Reactions from 1907 to 1985  
NEWS 11 Oct 24 BEILSTEIN adds new search fields  
NEWS 12 Oct 24 Nutraceuticals International (NUTRACEUT) now available on STN  
NEWS 13 Nov 18 DKILIT has been renamed APOLLIT  
NEWS 14 Nov 25 More calculated properties added to REGISTRY  
NEWS 15 Dec 04 CSA files on STN  
NEWS 16 Dec 17 PCTFULL now covers WP/PCT Applications from 1978 to date  
NEWS 17 Dec 17 TOXCENTER enhanced with additional content  
NEWS 18 Dec 17 Adis Clinical Trials Insight now available on STN  
NEWS 19 Jan 29 Simultaneous left and right truncation added to COMPENDEX,  
ENERGY, INSPEC  
NEWS 20 Feb 13 CANCERLIT is no longer being updated  
NEWS 21 Feb 24 METADEX enhancements  
NEWS 22 Feb 24 PCTGEN now available on STN  
NEWS 23 Feb 24 TEMA now available on STN  
NEWS 24 Feb 26 NTIS now allows simultaneous left and right truncation  
NEWS 25 Feb 26 PCTFULL now contains images  
NEWS 26 Mar 04 SDI PACKAGE for monthly delivery of multifile SDI results  
NEWS 27 Mar 20 EVENTLINE will be removed from STN  
NEWS 28 Mar 24 PATDPAFULL now available on STN  
NEWS 29 Mar 24 Additional information for trade-named substances without  
structures available in REGISTRY  
NEWS 30 Apr 11 Display formats in DGENE enhanced  
NEWS 31 Apr 14 MEDLINE Reload  
NEWS 32 Apr 17 Polymer searching in REGISTRY enhanced  
NEWS 33 Apr 21 Indexing from 1947 to 1956 being added to records in CA/CAPLUS  
NEWS 34 Apr 21 New current-awareness alert (SDI) frequency in  
WPIDS/WPINDEX/WPIX  
NEWS 35 Apr 28 RDISCLOSURE now available on STN  
NEWS 36 May 05 Pharmacokinetic information and systematic chemical names  
added to PHAR  
  
NEWS EXPRESS April 4 CURRENT WINDOWS VERSION IS V6.01a, CURRENT  
MACINTOSH VERSION IS V6.0b(ENG) AND V6.0Jb(JP),  
AND CURRENT DISCOVER FILE IS DATED 01 APRIL 2003  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 14:49:07 ON 15 MAY 2003

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'BIOSIS' ENTERED AT 14:49:30 ON 15 MAY 2003  
COPYRIGHT (C) 2003 BIOLOGICAL ABSTRACTS INC. (R)

FILE 'MEDLINE' ENTERED AT 14:49:30 ON 15 MAY 2003

FILE 'AGRICOLA' ENTERED AT 14:49:30 ON 15 MAY 2003

FILE 'EMBASE' ENTERED AT 14:49:30 ON 15 MAY 2003  
COPYRIGHT (C) 2003 Elsevier Science B.V. All rights reserved.

FILE 'CABA' ENTERED AT 14:49:30 ON 15 MAY 2003  
COPYRIGHT (C) 2003 CAB INTERNATIONAL (CABI)

FILE 'WPIDS' ENTERED AT 14:49:30 ON 15 MAY 2003  
COPYRIGHT (C) 2003 THOMSON DERWENT

FILE 'JAPIO' ENTERED AT 14:49:30 ON 15 MAY 2003  
COPYRIGHT (C) 2003 Japanese Patent Office (JPO)- JAPIO

FILE 'BIOTECHDS' ENTERED AT 14:49:30 ON 15 MAY 2003  
COPYRIGHT (C) 2003 THOMSON DERWENT AND INSTITUTE FOR SCIENTIFIC INFORMATION

FILE 'LIFESCI' ENTERED AT 14:49:30 ON 15 MAY 2003  
COPYRIGHT (C) 2003 Cambridge Scientific Abstracts (CSA)

FILE 'CAPLUS' ENTERED AT 14:49:30 ON 15 MAY 2003  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

=> e mansfield linda s/au

E1	4	MANSFIELD LEE A/AU
E2	1	MANSFIELD LINDA/AU
E3	29 -->	MANSFIELD LINDA S/AU
E4	1	MANSFIELD LORD/AU
E5	1	MANSFIELD LOUISE/AU
E6	2	MANSFIELD LUCIELLE/AU
E7	7	MANSFIELD LUCIELLE P/AU
E8	3	MANSFIELD LYNDON/AU
E9	9	MANSFIELD LYNDON E/AU
E10	76	MANSFIELD M/AU
E11	52	MANSFIELD M A/AU
E12	4	MANSFIELD M C/AU

=> s e2-e3  
L1 30 ("MANSFIELD LINDA"/AU OR "MANSFIELD LINDA S"/AU)

=> e mansfield l s/au  
E1 32 MANSFIELD L P/AU  
E2 6 MANSFIELD L R/AU  
E3 139 --> MANSFIELD L S/AU  
E4 24 MANSFIELD L W/AU  
E5 4 MANSFIELD LEE A/AU  
E6 1 MANSFIELD LINDA/AU  
E7 29 MANSFIELD LINDA S/AU  
E8 1 MANSFIELD LORD/AU  
E9 1 MANSFIELD LOUISE/AU  
E10 2 MANSFIELD LUCIELLE/AU  
E11 7 MANSFIELD LUCIELLE P/AU  
E12 3 MANSFIELD LYNDON/AU

=> s e3  
L2 139 "MANSFIELD L S"/AU

=> e rossano mary g/au  
E1 4 ROSSANO M J/AU  
E2 1 ROSSANO MARILENA/AU  
E3 9 --> ROSSANO MARY G/AU  
E4 56 ROSSANO P/AU  
E5 1 ROSSANO P G/AU  
E6 23 ROSSANO R/AU  
E7 1 ROSSANO R M/AU  
E8 3 ROSSANO R W/AU  
E9 1 ROSSANO ROBERT W/AU  
E10 7 ROSSANO ROCCO/AU  
E11 1 ROSSANO ROMAN MIGUEL/AU  
E12 13 ROSSANO S/AU

=> s e3  
L3 9 "ROSSANO MARY G"/AU

=> e rossano m g/au  
E1 4 ROSSANO M/AU  
E2 24 ROSSANO M A/AU  
E3 25 --> ROSSANO M G/AU  
E4 4 ROSSANO M J/AU  
E5 1 ROSSANO MARILENA/AU  
E6 9 ROSSANO MARY G/AU  
E7 56 ROSSANO P/AU  
E8 1 ROSSANO P G/AU  
E9 23 ROSSANO R/AU  
E10 1 ROSSANO R M/AU  
E11 3 ROSSANO R W/AU  
E12 1 ROSSANO ROBERT W/AU

=> s e3  
L4 25 "ROSSANO M G"/AU

=> e murphy alice j/au  
E1 2 MURPHY ALFRED M/AU  
E2 1 MURPHY ALICE/AU  
E3 14 --> MURPHY ALICE J/AU  
E4 2 MURPHY ALICE M/AU  
E5 5 MURPHY ALISON/AU  
E6 1 MURPHY ALISON A/AU  
E7 2 MURPHY ALLAN H/AU

E8	1	MURPHY ALLAN JAMES/AU
E9	1	MURPHY ALLAN R/AU
E10	1	MURPHY ALLEN MILTON JR/AU
E11	1	MURPHY ALLEYNE/AU
E12	3	MURPHY ALLISON R/AU

```
=> s e2 or e3
L5      15 "MURPHY ALICE"/AU OR "MURPHY ALICE J"/AU
```

```
=> e murphy a j/au
E1      24    MURPHY A I/AU
E2       2    MURPHY A I JR/AU
E3     337 --> MURPHY A J/AU
E4      19    MURPHY A J M/AU
E5       1    MURPHY A JR/AU
E6      10    MURPHY A K/AU
E7      77    MURPHY A L/AU
E8       3    MURPHY A LOLA/AU
E9       1    MURPHY A LYND/AU
E10     385    MURPHY A M/AU
E11     14    MURPHY A M C/AU
E12      5    MURPHY A M L/AU
```

```
=> s e3
L6     337 "MURPHY A J"/AU
```

```
=> e vtable ruth a/au
E1       6    VRABLE R G/AU
E2       2    VRABLE RUTH/AU
E3       9 --> VRABLE RUTH A/AU
E4       2    VRABLEC J J/AU
E5       1    VRABLEC J W/AU
E6      16    VRABLIC A S/AU
E7       5    VRABLIC ANGELICA S/AU
E8       1    VRABLIC ANGELICA SUAUA/AU
E9       1    VRABLIC DOROTHY M/AU
E10     50    VRABLIC J/AU
E11      2    VRABLIC M/AU
E12      3    VRABLIC O/AU
```

```
=> s e2 or e3
L7      11 "VRABLE RUTH"/AU OR "VRABLE RUTH A"/AU
```

```
=> e vtable r a/au
E1       7    VRABLE JOHN B/AU
E2      10    VRABLE R/AU
E3     13 --> VRABLE R A/AU
E4       6    VRABLE R G/AU
E5       2    VRABLE RUTH/AU
E6       9    VRABLE RUTH A/AU
E7       2    VRABLEC J J/AU
E8       1    VRABLEC J W/AU
E9      16    VRABLIC A S/AU
E10      5    VRABLIC ANGELICA S/AU
E11      1    VRABLIC ANGELICA SUAUA/AU
E12      1    VRABLIC DOROTHY M/AU
```

```
=> s e2 or e3
L8     23 "VRABLE R"/AU OR "VRABLE R A"/AU
```

```
=> s 11-18
L9    496 (L1 OR L2 OR L3 OR L4 OR L5 OR L6 OR L7 OR L8)
```

DT Article  
 LA English  
 AB Objective-To determine apparent seroprevalence of **antibodies** against **Sarcocystis neurona** in a population of domestic cats previously tested for **antibodies** against *Toxoplasma gondii*. Design-Cross-sectional study. Sample Population-Serum from 196 domestic cats. Procedure-Banked serum samples submitted to the Michigan State University Animal Health Diagnostic Laboratory for *T gondii* diagnostic testing were tested for **antibodies** against *S neurona* by use of an indirect fluorescent **antibody** (IFA) test and a western blot test. Submission records were analyzed to determine descriptive statistics and test for associations between positive results of a test for *S neurona* and other variables in the data set. Results-10 of 196 (5%) samples yielded positive results for **antibodies** against *S neurona* by use of western blot analysis, whereas 27 samples yielded positive results by use of the IFA. No association was found between *S neurona* western blot test results and *T gondii* test results, age, sex, or the reason for *T gondii* testing. The *S neurona* IFA titer was positively and significantly associated with positive results of western blot analysis. Conclusions and Clinical Relevance-Domestic cats are not likely to play a substantial role as intermediate hosts in the natural life cycle of *S neurona*. Results indicate that natural infection of domestic cats may occur, and small animal practitioners should be aware of this fact when evaluating cats with neurologic disease. The *S neurona* IFA test had lower specificity than western blot analysis.

L14 ANSWER 5 OF 12 WPIDS (C) 2003 THOMSON DERWENT DUPLICATE 3  
 AN 2001-218486 [22] WPIDS  
 CR 2000-571969 [49]  
 DNC C2001-065294  
 TI Vaccinating equids against protozoal **Sarcocystis neurona** infections using unique antigens.  
 DC B04 C06 D16  
 IN **MANSFIELD, L S; MURPHY, A J; ROSSANO, M G; VRABLE, R A**  
 PA (UNMS) UNIV MICHIGAN STATE  
 CYC 88  
 PI WO 2001015708 A1 20010308 (200122)\* EN 54p  
 RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
 NL OA PT SD SE SL SZ TZ UG ZW  
 W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB  
 GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS LT LU  
 LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR  
 TT UA UG UZ VN YU ZA ZW  
 AU 2000071087 A 20010326 (200137)  
 EP 1207889 A1 20020529 (200243) EN  
 R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
 RO SE SI  
 ADT WO 2001015708 A1 WO 2000-US24221 20000831; AU 2000071087 A AU 2000-71087  
 20000831; EP 1207889 A1 EP 2000-959829 20000831, WO 2000-US24221 20000831  
 FDT AU 2000071087 A Based on WO 200115708; EP 1207889 A1 Based on WO 200115708  
 PRAI US 2000-513086 20000224; US 1999-152193P 19990902  
 AB WO 200115708 A UPAB: 20020709  
 NOVELTY - Vaccinating equids against **Sarcocystis neurona** infections using polypeptide groups of unique 16 (+4) or 30 (+4) antigens of *S. neurona*, is new.  
 DETAILED DESCRIPTION - INDEPENDENT CLAIMS are included for the following:  
 (a) a vaccine (I) for providing passive immunity to **Sarcocystis neurona** infection, comprising **antibodies** against at least one group of a unique 16 (+4) or 30 (+4) antigen of *S. neurona*;  
 (b) a vaccine (II) for active immunization of an equid against a *S.*

neurona infection, comprising at least one group of a unique 16 (+4) or 30 (+4) antigen of *S. neurona*;

(c) a vaccine (III) for protecting an equid from *S. neurona* infection comprising a DNA that encodes at least 1 group of a 16 (+4) kDa antigen and/or a 30 (+4) kDa antigen of *S. neurona*;

(d) a method (IV) for vaccinating an equid against a *S. neurona* infection, comprising:

(1) providing a recombinant antigen of *S. neurona* produced from a recombinant microorganism culture (the microorganism contains a DNA that encodes at least one group of a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona*; and

(2) vaccinating the equid;

(e) a method (V) for vaccinating an equid against a *S. neurona* infection, comprising:

(1) providing a DNA in a carrier solution, a plasmid which encodes at least 1 group of a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of ***Sarcocystis neurona***; and

(2) vaccinating the equid with the DNA in the carrier solution;

(f) a method (VI) of providing passive immunity to a *S. neurona* infection in a equid, comprising:

(1) providing **antibodies** against at least 1 group of a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona* (the **antibodies** may be monoclonal or polyclonal); and

(2) inoculating the equid;

(g) a method (VII) for producing a polypeptide, comprising:

(1) providing a microorganism in a culture containing a DNA encoding a fusion polypeptide comprising at least 1 group of a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona* and a polypeptide that facilitates isolation of the fusion polypeptide;

(2) culturing the microorganism in a culture to produce the fusion polypeptide; and

(3) isolating the fusion polypeptide;

(h) a method (VIII) for producing an **antibody** comprising:

(1) providing a microorganism in a culture containing DNA encoding a fusion polypeptide comprising at least 1 group of a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona* and a polypeptide that facilitates isolation of the fusion polypeptide;

(2) culturing the microorganisms in a culture to produce the fusion polypeptide;

(3) isolating the fusion polypeptide;

(4) producing the **antibody** from the polypeptide;

(i) a monoclonal **antibody** (IX) that selectively binds to a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen;

(j) an isolated DNA (X) encoding a monoclonal **antibody** that selectively binds to a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen;

(k) a bacterial clone (XI) containing a plasmid comprising a DNA encoding a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona*;

(l) a vaccine (XII) for an equid comprising an isolated recombinant protein encoded by a cDNA produced from mRNA of *S. neurona* encoding a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen;

(m) a vaccine (XIII) for an equid comprising a recombinant virus vector containing DNA encoding a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona*;

(n) a DNA vaccine (XIV) for an equid comprising a plasmid containing DNA encoding a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona*; and

(o) a method (XV) for protecting an equid against *S. neurona* which comprises providing a vaccine that when injected into the equid causes the equid to produce antibodies against a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona* (the antibodies prevent infection by the *Sarcocystis neurona*).

ACTIVITY - Antiparasitic.

No biological data given.

(stratified by the state's opossum (*Didelphis virginiana*) population and the number of equids on each operation) was selected. Ninety-eight equine-operation owners agreed to participate, and blood collection occurred from late March through October of 1997. Data regarding the 98 farms' feeding and management practices were collected, as well as descriptive data for each of the 1121 individual horses. Serum samples were tested for **antibodies** to *S. neurona* using a Western blot test. The true seroprevalence of **antibodies** specific to *S. neurona* was estimated to be 60%. Chi-square analysis showed that seroprevalence was lowest in the colder parts of the state that had the fewest opossums ( $P < 0.0001$ ). In two multivariable logistic-regression analyses with random effects grouped by herd, age and exposure to pasture were associated with increased odds of seropositivity, and feeding of sweet feed (grains mixed with molasses) was associated with decreased odds of testing positive. No association was found between farm size, animal gender, hay types, horse-housing types or exposure to natural surface water and seropositivity.

L14 ANSWER 8 OF 12 WPIDS (C) 2003 THOMSON DERWENT DUPLICATE 5  
 AN 2000-571969 [53] WPIDS  
 CR 2001-218486 [22]  
 DNN N2000-423167 DNC C2000-170452  
 TI Detection of **Sarcocystis neurona**, which causes equine protozoal myeloencephalitis, in horse serum and cerebrospinal fluid comprises identifying a specific **antibody**-antigen complex via an immunoassay.  
 DC B04 C07 D16 S03  
 IN **MANSFIELD, L S; MURPHY, A J; ROSSANO, M G; VRABLE, R A**  
 PA (UNMS) UNIV MICHIGAN STATE  
 CYC 87  
 PI WO 2000049049 A1 20000824 (200053)\* EN 64p  
 RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
 OA PT SD SE SL SZ TZ UG ZW  
 W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB  
 GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS LT LU  
 LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR  
 TT UA UG UZ VN YU ZA ZW  
 AU 2000034982 A 20000904 (200103)  
 US 6344337 B1 20020205 (200211)  
 ADT WO 2000049049 A1 WO 2000-US4379 20000218; AU 2000034982 A AU 2000-34982  
 20000218; US 6344337 B1 Provisional US 1999-120831P 19990219, Provisional  
 US 1999-152193P 19990902, US 2000-506630 20000218  
 FDT AU 2000034982 A Based on WO 200049049  
 PRAI US 1999-152193P 19990902; US 1999-120831P 19990219; US 2000-506630  
 20000218  
 AB WO 200049049 A UPAB: 20020215  
 NOVELTY - Detection of **Sarcocystis neurona** in horses by identifying a specific **antibody**-antigen complex via an immunoassay is new.  
 DETAILED DESCRIPTION - Detection of **Sarcocystis neurona** in an equine in an immunoassay is improved by reacting a biological sample from the horse suspected of harboring the *S. neurona* with an **antibody** (Ab) which is selective in binding to an identifying *S. neurona* antigen (Ag) to form an Ab-Ag complex.  
 INDEPENDENT CLAIMS are also included for the following:  
 (1) a kit for detecting *S. neurona* in a biological sample from an equine;  
 (2) monoclonal **antibodies** against 16 plus or minus 4 kDa or 30 plus or minus 4 kDa antigens of *S. neurona*; and  
 (3) isolated DNA sequences encoding the 16 plus or minus 4 kDa and 30 plus or minus 4 kDa antigens of *S. neurona*.  
 USE - The methods and **antibodies** are useful for detecting

S. neurona (claimed) which causes equine protozoal myeloencephalitis, a neurological disorder in horses.  
Dwg.0/0

L14 ANSWER 9 OF 12 WPIDS (C) 2003 THOMSON DERWENT DUPLICATE 6  
AN 2000-292877 [25] WPIDS  
DNN N2000-219631 DNC C2000-088472  
TI Immunoassay for equine protozoal myeloencephalitis in horses uses specific **antibodies** to proteins derived from **Sarcocystis neurona**.  
DC B04 C06 D16 S03  
IN **MANSFIELD, L S; MURPHY, A J; ROSSANO, M G**  
PA (UNMS) UNIV MICHIGAN STATE  
CYC 83  
PI WO 2000017640 A1 20000330 (200025)\* EN 26p  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ UG ZW  
W: AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB GE  
GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MD  
MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR TT UA  
UG UZ VN YU ZW  
AU 9954707 A 20000410 (200035)  
US 6153394 A 20001128 (200063)  
US 6489148 B1 20021203 (200301)  
ADT WO 2000017640 A1 WO 1999-US17961 19990809; AU 9954707 A AU 1999-54707  
19990809; US 6153394 A US 1998-156954 19980918; US 6489148 B1 Div ex US  
1998-156954 19980918, US 2000-569434 20000512  
FDT AU 9954707 A Based on WO 200017640; US 6489148 B1 Div ex US 6153394  
PRAI US 1998-156954 19980918; US 2000-569434 20000512  
AB WO 200017640 A UPAB: 20000524  
NOVELTY - An improved immunoassay for detecting **Sarcocystis neurona** infection in equines, comprises reacting the **Sarcocystis neurona** protein with a non-labeled **antibody** to proteins of other **Sarcocystis** species, before the immunoassay, which inhibits non-specific binding of the labeled **antibody**, during the immunoassay.  
DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following:  
(1) a method for the detection of disease caused by **Sarcocystis** neurons in equines which comprises:  
(a) isolating fluid from the equine which can contain parasite induced **antibodies** to **Sarcocystis neurona** proteins, indicating the presence of the **Sarcocystis neurona**;  
(b) reacting the fluid with at least one identifying antigen of the **Sarcocystis** neurons protein bound on a substrate, where the substrate has been blocked with **antibodies** to **Sarcocystis** sp. other than **Sarcocystis** neurons, so that **antibodies** to **Sarcocystis neurona** antigen in the fluid are bound to the identifying antigen;  
and  
(c) detecting the **antibodies** bound to the antigen;  
(2) a kit for the detection of disease caused by **Sarcocystis neurona** comprising in separate containers:  
(a) an identifying **antibody** able to specifically bind a **Sarcocystis neurona** protein; and  
(b) a non-labeled **antibody** which is specific for a second protein of a **Sarcocystis** sp. other than **Sarcocystis neurona**; and  
(3) a kit for the detection of disease caused by **Sarcocystis neurona** in equines comprising:  
(a) a substrate with at least one identifying antigen to the **Sarcocystis neurona** bound on a surface of the substrate;  
(b) **antibody** to a **Sarcocystis** sp. other than



**Sarcocystis neurona**; and

(c) at least one reagent for the detection of an **antibody** in a fluid of the equine which binds to the antigen of **Sarcocystis neurona**.

USE - The methods and kits are used to detect **antibodies** to proteins of **Sarcocystis neurona**, in an equine, (claimed), which causes myeloencephalitis in the equine.

ADVANTAGE - The method uses a non-labeled **antibody** to proteins of other **Sarcocystis** species to inhibit the non-specific binding of the labeled **antibody**, improving the accuracy of the assay.

Dwg.0/2

L14 ANSWER 10 OF 12 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
AN 2001:258224 BIOSIS  
DN PREV200100258224  
TI Immunoassay for equine protozoal myeloencephalitis in horses.  
AU **Mansfield, Linda S. (1); Murphy, Alice J.;  
Rossano, Mary G.**  
CS (1) Bath, MI USA  
ASSIGNEE: Board of Trustees operating Michigan State University  
PI US 6153394 November 28, 2000  
SO Official Gazette of the United States Patent and Trademark Office Patents,  
(Nov. 28, 2000) Vol. 1240, No. 4, pp. No Pagination. e-file.  
ISSN: 0098-1133.  
DT Patent  
LA English  
AB An immunoassay for **Sarcocystis neurona**  
**antibodies** in equines is described. The immunoassay uses blocking  
of **Sarcocystis** antigens by **antibodies** to **Sarcocystis** sp. other  
than **Sarcocystis neurona** in connection with the  
immunoassay.

L14 ANSWER 11 OF 12 BIOTECHDS COPYRIGHT 2003 THOMSON DERWENT AND ISI  
AN 2000-14646 BIOTECHDS  
TI Detection of **Sarcocystis neurona**, which causes horse  
protozoan myeloencephalitis, in horse serum and cerebrospinal fluid  
comprises identifying a specific **antibody**-antigen complex via  
an immunoassay;  
with use of monoclonal **antibodies**  
AU **Mansfield L S; Rossano M G; Murphy A J;  
Vrable R A**  
PA Univ.Michigan-State  
LO East Lansing, MI, USA.  
PI WO 2000049049 24 Aug 2000  
AI WO 2000-US4379 18 Feb 2000  
PRAI US 990152193 2 Sep 1999; US 1999-120831 19 Feb 1999  
DT Patent  
LA English  
OS WPI: 2000-571969 [53]  
AB Detection of **Sarcocystis neurona**, which causes equine  
protozoan myeloencephalitis in horses by identifying a specific  
**antibody**-antigen complex via an immunoassay, is claimed. Also  
claimed are: a kit for detecting **S. neurona** in a biological sample from a  
horse; monoclonal **antibodies** against antigens of **S. neurona**;  
and isolated DNA sequences encoding the antigens of **S. neurona**. The  
methods and **antibodies** are useful for detecting **S. neurona**  
which causes horse protozoan myeloencephalitis, a neurological disorder  
in horses. The labelled **antibody** against the antigen or the  
**antibody** in the **antibody**-antigen complex is provided  
for the detecting. The label is chosen from alkaline phosphatase  
(EC-3.1.3.1), horseradish peroxidase (EC-1.11.1.7), fluorescent  
compounds, luminescent compounds, colloidal gold and magnetic particles.  
The label is preferably biotin, which is reacted with peroxidase

L3           9 S E3  
             E ROSSANO MARY G/AU  
 L4           25 S E3  
             E ROSSANO M G/AU  
 L5           15 S E2 OR E3  
             E MURPHY ALICE J/AU  
 L6           337 S E3  
             E VRABLE RUTH A/AU  
 L7           11 S E2 OR E3  
             E VRABLE R A/AU  
 L8           23 S E2 OR E3  
 L9           496 S L1-L8  
 L10          39 S L9 AND SARCOCYSTIS NEURONA  
 L11          1 S L10 AND (16 KD OR 30 KD)  
 L12          51 S L9 AND ANTIBOD?  
 L13          26 S L10 AND ANTIBOD?  
 L14          12 DUP REM L13 (14 DUPLICATES REMOVED)

=> dup rem l10

PROCESSING COMPLETED FOR L10

L15           15 DUP REM L10 (24 DUPLICATES REMOVED)

=> d bib ab 1-15

L15 ANSWER 1 OF 15 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE  
1

AN 2003:159207 BIOSIS

DN PREV200300159207

TI A herd-level analysis of risk factors for antibodies to  
**Sarcocystis neurona** in Michigan equids.

AU **Rossano, M. G.**; Kaneene, J. B. (1); Marteniuk, J. V.; Banks, B.  
D.; Schott, H. C., II; **Mansfield, L. S.**

CS (1) College of Veterinary Medicine, Population Medicine Center, A-109  
Veterinary Medical Center, Michigan State University, East Lansing, MI,  
48824-1314, USA: kaneene@cvm.msu.edu USA

SO Preventive Veterinary Medicine, (15 February 2003) Vol. 57, No. 1-2, pp.  
7-13. print.

ISSN: 0167-5877.

DT Article

LA English

AB Equine protozoal myeloencephalitis (EPM) is a neurological disease of horses and ponies caused by infection of the central nervous system with the protozoan parasite **Sarcocystis neurona**. A herd-level analysis of a cross-sectional study of serum antibodies to *S. neurona* in Michigan equids was conducted, using data collected in 1997 for study that included 1121 equids from 98 Michigan horse farms. Our objective was to identify specific herd-level risk factors associated with seropositivity. We tested associations between herd seroprevalence and various farm-management practices (including feed-storage methods and wildlife control). Multivariable models were developed for three strata based on relative opossum abundance (opossum districts). Herd seroprevalence ranged from 0 to 100% (median = 57%); No risk factor was significantly associated with herd seroprevalence at  $P \leq 0.05$  in all opossum districts. Our results suggest that equids living in areas with large opossum populations might be infected with *S. neurona* from multiple sources.

L15 ANSWER 2 OF 15 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

AN 2003:67219 BIOSIS

DN PREV200300067219

TI Immunoassay for equine protozoal myeloencephalitis in horses.

AU **Mansfield, Linda S.**; **Murphy, Alice J.** (1);

fluorescent antibody (IFA) test and a western blot test. Submission records were analyzed to determine descriptive statistics and test for associations between positive results of a test for *S. neurona* and other variables in the data set. Results-10 of 196 (5%) samples yielded positive results for antibodies against *S. neurona* by use of western blot analysis, whereas 27 samples yielded positive results by use of the IFA. No association was found between *S. neurona* western blot test results and *T. gondii* test results, age, sex, or the reason for *T. gondii* testing. The *S. neurona* IFA titer was positively and significantly associated with positive results of western blot analysis. Conclusions and Clinical Relevance-Domestic cats are not likely to play a substantial role as intermediate hosts in the natural life cycle of *S. neurona*. Results indicate that natural infection of domestic cats may occur, and small animal practitioners should be aware of this fact when evaluating cats with neurologic disease. The *S. neurona* IFA test had lower specificity than western blot analysis.

L15 ANSWER 5 OF 15 WPIDS (C) 2003 THOMSON DERWENT DUPLICATE 3  
 AN 2001-218486 [22] WPIDS  
 CR 2000-571969 [49]  
 DNC C2001-065294  
 TI Vaccinating equids against protozoal **Sarcocystis neurona** infections using unique antigens.  
 DC B04 C06 D16  
 IN MANSFIELD, L S; MURPHY, A J; ROSSANO, M G; VRABLE, R A  
 PA (UNMS) UNIV MICHIGAN STATE  
 CYC 88  
 PI WO 2001015708 A1 20010308 (200122)\* EN 54p  
 RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ NL OA PT SD SE SL SZ TZ UG ZW  
 W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR TT UA UG UZ VN YU ZA ZW  
 AU 2000071087 A 20010326 (200137)  
 EP 1207889 A1 20020529 (200243) EN  
 R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT RO SE SI  
 ADT WO 2001015708 A1 WO 2000-US24221 20000831; AU 2000071087 A AU 2000-71087 20000831; EP 1207889 A1 EP 2000-959829 20000831, WO 2000-US24221 20000831  
 FDT AU 2000071087 A Based on WO 200115708; EP 1207889 A1 Based on WO 200115708  
 PRAI US 2000-513086 20000224; US 1999-152193P 19990902  
 AB WO 200115708 A UPAB: 20020709  
 NOVELTY - Vaccinating equids against **Sarcocystis neurona** infections using polypeptide groups of unique 16 (+4) or 30 (+4) antigens of *S. neurona*, is new.  
 DETAILED DESCRIPTION - INDEPENDENT CLAIMS are included for the following:  
 (a) a vaccine (I) for providing passive immunity to **Sarcocystis neurona** infection, comprising antibodies against at least one group of a unique 16 (+4) or 30 (+4) antigen of *S. neurona*;  
 (b) a vaccine (II) for active immunization of an equid against a *S. neurona* infection, comprising at least one group of a unique 16 (+4) or 30 (+4) antigen of *S. neurona*;  
 (c) a vaccine (III) for protecting an equid from *S. neurona* infection comprising a DNA that encodes at least 1 group of a 16 (+4) kDa antigen and/or a 30 (+4) kDa antigen of *S. neurona*;  
 (d) a method (IV) for vaccinating an equid against a *S. neurona* infection, comprising:  
 (1) providing a recombinant antigen of *S. neurona* produced from a recombinant microorganism culture (the microorganism contains a DNA that

encodes at least one group of a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona*; and

(2) vaccinating the equid;

(e) a method (V) for vaccinating an equid against a *S. neurona* infection, comprising:

(1) providing a DNA in a carrier solution, a plasmid which encodes at least 1 group of a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of

***Sarcocystis neurona***; and

(2) vaccinating the equid with the DNA in the carrier solution;

(f) a method (VI) of providing passive immunity to a *S. neurona* infection in a equid, comprising:

(1) providing antibodies against at least 1 group of a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona* (the antibodies may be monoclonal or polyclonal); and

(2) inoculating the equid;

(g) a method (VII) for producing a polypeptide, comprising:

(1) providing a microorganism in a culture containing a DNA encoding a fusion polypeptide comprising at least 1 group of a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona* and a polypeptide that facilitates isolation of the fusion polypeptide;

(2) culturing the microorganism in a culture to produce the fusion polypeptide; and

(3) isolating the fusion polypeptide;

(h) a method (VIII) for producing an antibody comprising:

(1) providing a microorganism in a culture containing DNA encoding a fusion polypeptide comprising at least 1 group of a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona* and a polypeptide that facilitates isolation of the fusion polypeptide;

(2) culturing the microorganisms in a culture to produce the fusion polypeptide;

(3) isolating the fusion polypeptide;

(4) producing the antibody from the polypeptide;

(i) a monoclonal antibody (IX) that selectively binds to a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen;

(j) an isolated DNA (X) encoding a monoclonal antibody that selectively binds to a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen;

(k) a bacterial clone (XI) containing a plasmid comprising a DNA encoding a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona*;

(l) a vaccine (XII) for an equid comprising an isolated recombinant protein encoded by a cDNA produced from mRNA of *S. neurona* encoding a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen;

(m) a vaccine (XIII) for an equid comprising a recombinant virus vector containing DNA encoding a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona*;

(n) a DNA vaccine (XIV) for an equid comprising a plasmid containing DNA encoding a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona*; and

(o) a method (XV) for protecting an equid against *S. neurona* which comprises providing a vaccine that when injected into the equid causes the equid to produce antibodies against a 16 (+4) kDa antigen and/or 30 (+4) kDa antigen of *S. neurona* (the antibodies prevent infection by the *Sarcocystis neurona*).

ACTIVITY - Antiparasitic.

No biological data given.

MECHANISM OF ACTION - Vaccine.

USE - The vaccines and methods are used for protecting equids against infections by the protozoan parasite *Sarcocystis neurona*.

Dwg.0/0

**Sarcocystis neurona** nucleic acid vaccine and  
recombinant vaccine

AU **Mansfield L S; Rossano M G; Murphy A J;  
Vrable R A**

PA Univ.Michigan-State

LO East Lansing, MI, USA.

PI WO 2001015708 8 Mar 2001

AI WO 2000-US24221 31 Aug 2000

PRAI US 2000-513086 24 Feb 2000; US 1999-152193 2 Sep 1999

DT Patent

LA English

OS WPI: 2001-218486 [22]

AB A method for vaccinating equids against **Sarcocystis neurona** infection is claimed. It involves using protein groups of unique 16(+4) or 30(+4) antigens of *S. neurona*. Also claimed are: a vaccine (I) for providing passive immunity to **Sarcocystis neurona** infection; a vaccine (II) for active immunization of an equid against a *S. neurona* infection; a vaccine (III) for protecting an equid from *S. neurona* infection; a method (IV or V) for vaccinating an equid against a *S. neurona* infection; a method (VI) of providing passive immunity to a *S. neurona* infection; a method (VII) for producing a protein (e.g. glutathione-transferase); a method (VIII) for producing an antibody; providing a microorganism in a culture containing DNA encoding a fusion protein; a monoclonal antibody (IX); an isolated DNA (X); a bacterial clone (XI); a vaccine (XII) for an equid containing an isolated recombinant protein; a vaccine (XIII or XIV) for an equid containing recombinant virus vector containing DNA; and a method (XV) for protecting an equid against *S. neurona*. The vaccines and methods are used for protecting equids against infection by the protozoon parasite *Sarcocystis neurona*. (54pp)

L15 ANSWER 7 OF 15 CABA COPYRIGHT 2003 CABI

AN 2001:140181 CABA

DN 20013139279

TI The effects of pyrantel tartrate on **Sarcocystis neurona**  
merozoite viability

AU Kruttlin, E. A.; Rossano, M. G.; Murphy, A. J.;  
Vrable, R. A.; Kaneene, J. B.; Schott, H. C., II; Mansfield,  
L. S.

CS Department of Large Animal Clinical Sciences, D201 Veterinary Medicine  
Center, College of Veterinary Medicine, Michigan State University, East  
Lansing, MI 48824, USA.

SO Veterinary Therapeutics, (2001) Vol. 2, No. 3, pp. 268-276. 21 ref.  
ISSN: 1528-3593

DT Journal

LA English

AB *S. neurona* is the etiologic agent of equine protozoal myeloencephalitis, a neurologic disease of horses. The present study was designed to test the hypothesis that pyrantel tartrate can kill *S. neurona* merozoites growing in equine dermal cell culture. *S. neurona* merozoites were exposed to a range of concentrations of pyrantel tartrate or sodium tartrate ranging from 0.001 to 0.01 M. Merozoites were then placed onto equine dermal cell cultures and incubated for 2 weeks to check for viability. At 1 and 2 weeks after inoculation, plaque counts were compared between treatments and, between treatments and controls. Merozoites exposed to concentrations of pyrantel tartrate higher than 0.0025 M (8.91x10<sup>-4</sup> g/ml) did not produce plaques in equine dermal cells, whereas those exposed to similar concentrations of the tartrate salt or medium alone produced significant numbers of plaques. These results that pyrantel tartrate has activity against *S. neurona* merozoites in vitro and suggest that it may have activity against the sporozoite stage of the parasite found in the equine gut.

L15 ANSWER 8 OF 15 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE

4

AN 2001:169884 BIOSIS

DN PREV200100169884

TI Comparison of *Sarcocystis neurona* isolates derived from horse neural tissue.

AU **Mansfield, L. S. (1)**; Schott, H. C., II; **Murphy, A. J.**

; **Rossano, M. G.**; Tanhauser, S. M.; Patterson, J. S.; Nelson, K.; Ewart, S. L.; Marteniuk, J. V.; Bowman, D. D.; Kaneene, J. B.

CS (1) Department of Large Animal Clinical Sciences, College of Veterinary Medicine, Michigan State University, East Lansing, MI, 48824: mansfie4@cvm.msu.edu USA

SO Veterinary Parasitology, (26 February, 2001) Vol. 95, No. 2-4, pp. 167-178. print.  
ISSN: 0304-4017.

DT Article

LA English

SL English

AB *Sarcocystis neurona* is a protozoan parasite that can cause neurological deficits in infected horses. The route of transmission is by fecal-oral transfer of sporocysts from opossums. However, the species identity and the lifecycle are not completely known. In this study, *Sarcocystis* merozoites from eight isolates obtained from Michigan horses were compared to *S. neurona* from a California horse (UCD1), *Sarcocystis* from a grackle (Cornell), and five *Sarcocystis* isolates from feral opossums from Michigan. Comparisons were made using several techniques. SDS-PAGE analysis with silver staining showed that *Sarcocystis* spp. from the eight horses appeared the same, but different from the grackle isolate. One Michigan horse isolate (MIH6) had two bands at 72 and 25 kDa that were more prominent than the UCD1 isolate and other Michigan horse isolates. Western blot analysis showed that merozoites of eight of eight equine-derived isolates, and the UCD1 *S. neurona* isolate had similar bands when developed with serum or CSF of an infected horse. Major bands were seen at 60, 44, 30, and 16 kDa. In the grackle (Cornell) isolate, bands were seen at 60, 44, 29, and 16 kDa. DNA from merozoites of each of the eight equine-derived isolates and the grackle-derived isolate produced a 334 bp PCR product (Tanhauser et al., 1999). Restriction fragment length polymorphism (RFLP) analysis of these horse isolates showed banding patterns characteristic for *S. neurona*. The grackle (Cornell) isolate had an RFLP banding pattern characteristic of other *S. falcatula* species. Finally, electron microscopy examining multiple merozoites of each of these eight horse isolates showed similar morphology, which differed from the grackle (Cornell) isolate. We conclude that the eight Michigan horse isolates are *S. neurona* species and the grackle isolate is an *S. falcatula* species.

L15 ANSWER 9 OF 15 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE

5

AN 2001:135932 BIOSIS

DN PREV200100135932

TI The seroprevalence of antibodies to *Sarcocystis neurona* in Michigan equids.

AU **Rossano, M. G.**; Kaneene, J. B. (1); Marteniuk, J. V.; Banks, B. D.; Schott, H. C., II; **Mansfield, L. S.**

CS (1) Population Medicine Center, College of Veterinary Medicine, A-109 Veterinary Medical Center, Michigan State University, East Lansing, MI, 48824-1314: kaneene@cvm.msu.edu USA

SO Preventive Veterinary Medicine, (29 January, 2001) Vol. 48, No. 2, pp. 113-128. print.  
ISSN: 0167-5877.

DT Article

LA English

SL English

AB A cross-sectional study of serum antibodies to **Sarcocystis neurona** (the etiologic agent of equine protozoal myeloencephalitis, EPM) was performed on Michigan equids. Our objectives were to determine the seroprevalence of antibodies to *S. neurona* in Michigan equids and to identify specific risk factors for seropositivity. A random, weighted sample of Michigan horse farms (stratified by the state's opossum (*Didelphis virginiana*) population and the number of equids on each operation) was selected. Ninety-eight equine-operation owners agreed to participate, and blood collection occurred from late March through October of 1997. Data regarding the 98 farms' feeding and management practices were collected, as well as descriptive data for each of the 1121 individual horses. Serum samples were tested for antibodies to *S. neurona* using a Western blot test. The true seroprevalence of antibodies specific to *S. neurona* was estimated to be 60%. Chi-square analysis showed that seroprevalence was lowest in the colder parts of the state that had the fewest opossums ( $P < 0.0001$ ). In two multivariable logistic-regression analyses with random effects grouped by herd, age and exposure to pasture were associated with increased odds of seropositivity, and feeding of sweet feed (grains mixed with molasses) was associated with decreased odds of testing positive. No association was found between farm size, animal gender, hay types, horse-housing types or exposure to natural surface water and seropositivity.

L15 ANSWER 10 OF 15 WPIDS (C) 2003 THOMSON DERWENT DUPLICATE 6

AN 2000-571969 [53] WPIDS

CR 2001-218486 [22]

DNN N2000-423167 DNC C2000-170452

TI Detection of **Sarcocystis neurona**, which causes equine protozoal myeloencephalitis, in horse serum and cerebrospinal fluid comprises identifying a specific antibody-antigen complex via an immunoassay.

DC B04 C07 D16 S03

IN MANSFIELD, L S; MURPHY, A J; ROSSANO, M G;  
VRABLE, R A

PA (UNMS) UNIV MICHIGAN STATE

CYC 87

PI WO 2000049049 A1 20000824 (200053)\* EN 64p

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ TZ UG ZW

W: AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB  
GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS LT LU  
LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR  
TT UA UG UZ VN YU ZA ZW

AU 2000034982 A 20000904 (200103)

US 6344337 B1 20020205 (200211)

ADT WO 2000049049 A1 WO 2000-US4379 20000218; AU 2000034982 A AU 2000-34982  
20000218; US 6344337 B1 Provisional US 1999-120831P 19990219, Provisional  
US 1999-152193P 19990902, US 2000-506630 20000218

FDT AU 2000034982 A Based on WO 200049049

PRAI US 1999-152193P 19990902; US 1999-120831P 19990219; US 2000-506630  
20000218

AB WO 200049049 A UPAB: 20020215

NOVELTY - Detection of **Sarcocystis neurona** in horses  
by identifying a specific antibody-antigen complex via an immunoassay is  
new.

DETAILED DESCRIPTION - Detection of **Sarcocystis neurona**  
**neurona** in an equine in an immunoassay is improved by reacting a  
biological sample from the horse suspected of harboring the *S. neurona*  
with an antibody (Ab) which is selective in binding to an identifying *S.*  
*neurona* antigen (Ag) to form an Ab-Ag complex.

INDEPENDENT CLAIMS are also included for the following:

(1) a kit for detecting *S. neurona* in a biological sample from an  
equine;

(2) monoclonal antibodies against 16 plus or minus 4 kDa or 30 plus or minus 4 kDa antigens of *S. neurona*; and  
(3) isolated DNA sequences encoding the 16 plus or minus 4 kDa and 30 plus or minus 4 kDa antigens of *S. neurona*.

USE - The methods and antibodies are useful for detecting *S. neurona* (claimed) which causes equine protozoal myeloencephalitis, a neurological disorder in horses.

Dwg.0/0

L15 ANSWER 11 OF 15 WPIDS (C) 2003 THOMSON DERWENT DUPLICATE 7  
AN 2000-292877 [25] WPIDS  
DNN N2000-219631 DNC C2000-088472  
TI Immunoassay for equine protozoal myeloencephalitis in horses uses specific antibodies to proteins derived from **Sarcocystis neurona**

DC B04 C06 D16 S03  
IN **MANSFIELD, L S; MURPHY, A J; ROSSANO, M G**  
PA (UNMS) UNIV MICHIGAN STATE  
CYC 83

PI WO 2000017640 A1 20000330 (200025)\* EN 26p  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL  
OA PT SD SE SL SZ UG ZW  
W: AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB GE  
GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MD  
MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR TT UA  
UG UZ VN YU ZW

AU 9954707 A 20000410 (200035)  
US 6153394 A 20001128 (200063)  
US 6489148 B1 20021203 (200301)

ADT WO 2000017640 A1 WO 1999-US17961 19990809; AU 9954707 A AU 1999-54707  
19990809; US 6153394 A US 1998-156954 19980918; US 6489148 B1 Div ex US  
1998-156954 19980918, US 2000-569434 20000512

FDT AU 9954707 A Based on WO 200017640; US 6489148 B1 Div ex US 6153394

PRAI US 1998-156954 19980918; US 2000-569434 20000512

AB WO 200017640 A UPAB: 20000524

NOVELTY - An improved immunoassay for detecting **Sarcocystis neurona** infection in equines, comprises reacting the **Sarcocystis neurona** protein with a non-labeled antibody to proteins of other **Sarcocystis** species, before the immunoassay, which inhibits non-specific binding of the labeled antibody, during the immunoassay.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following:

(1) a method for the detection of disease caused by **Sarcocystis** neurons in equines which comprises:

(a) isolating fluid from the equine which can contain parasite induced antibodies to **Sarcocystis neurona** proteins, indicating the presence of the **Sarcocystis neurona**;

(b) reacting the fluid with at least one identifying antigen of the **Sarcocystis** neurons protein bound on a substrate, where the substrate has been blocked with antibodies to **Sarcocystis** sp. other than **Sarcocystis** neurons, so that antibodies to **Sarcocystis neurona** antigen in the fluid are bound to the identifying antigen; and

(c) detecting the antibodies bound to the antigen;

(2) a kit for the detection of disease caused by **Sarcocystis neurona** comprising in separate containers:

(a) an identifying antibody able to specifically bind a **Sarcocystis neurona** protein; and

(b) a non-labeled antibody which is specific for a second protein of a **Sarcocystis** sp. other than **Sarcocystis neurona**; and

(3) a kit for the detection of disease caused by **Sarcocystis neurona** in equines comprising:

(a) a substrate with at least one identifying antigen to the



**Sarcocystis neurona** bound on a surface of the substrate;

(b) antibody to a **Sarcocystis** sp. other than **Sarcocystis neurona**; and

(c) at least one reagent for the detection of an antibody in a fluid of the equine which binds to the antigen of **Sarcocystis neurona**.

USE - The methods and kits are used to detect antibodies to proteins of **Sarcocystis neurona**, in an equine, (claimed), which causes myeloencephalitis in the equine.

ADVANTAGE - The method uses a non-labeled antibody to proteins of other **Sarcocystis** species to inhibit the non-specific binding of the labeled antibody, improving the accuracy of the assay.

Dwg.0/2

L15 ANSWER 12 OF 15 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

AN 2001:258224 BIOSIS

DN PREV200100258224

TI Immunoassay for equine protozoal myeloencephalitis in horses.

AU **Mansfield, Linda S. (1); Murphy, Alice J.;**

**Rossano, Mary G.**

CS (1) Bath, MI USA

ASSIGNEE: Board of Trustees operating Michigan State University

PI US 6153394 November 28, 2000

SO Official Gazette of the United States Patent and Trademark Office Patents, (Nov. 28, 2000) Vol. 1240, No. 4, pp. No Pagination. e-file.

ISSN: 0098-1133.

DT Patent

LA English

AB An immunoassay for **Sarcocystis neurona** antibodies in equines is described. The immunoassay uses blocking of **Sarcocystis** antigens by antibodies to **Sarcocystis** sp. other than **Sarcocystis neurona** in connection with the immunoassay.

L15 ANSWER 13 OF 15 BIOTECHDS COPYRIGHT 2003 THOMSON DERWENT AND ISI

AN 2000-14646 BIOTECHDS

TI Detection of **Sarcocystis neurona**, which causes horse protozoan myeloencephalitis, in horse serum and cerebrospinal fluid comprises identifying a specific antibody-antigen complex via an immunoassay;

with use of monoclonal antibodies

AU **Mansfield L S; Rossano M G; Murphy A J;**

**Vrable R A**

PA Univ.Michigan-State

LO East Lansing, MI, USA.

PI WO 2000049049 24 Aug 2000

AI WO 2000-US4379 18 Feb 2000

PRAI US 990152193 2 Sep 1999; US 1999-120831 19 Feb 1999

DT Patent

LA English

OS WPI: 2000-571969 [53]

AB Detection of **Sarcocystis neurona**, which causes equine protozoan myeloencephalitis in horses by identifying a specific antibody-antigen complex via an immunoassay, is claimed. Also claimed are: a kit for detecting **S. neurona** in a biological sample from a horse; monoclonal antibodies against antigens of **s. neurona**; and isolated DNA sequences encoding the antigens of **S. neurona**. The methods and antibodies are useful for detecting **S. neurona** which causes horse protozoan myeloencephalitis, a neurological disorder in horses. The labelled antibody against the antigen or the antibody in the antibody-antigen complex is provided for the detecting. The label is chosen from alkaline phosphatase (EC-3.1.3.1), horseradish peroxidase (EC-1.11.1.7), fluorescent compounds, luminescent compounds, colloidal gold and magnetic particles. The label is preferably biotin, which is

LA English

SL English

AB **Sarcocystis neurona** is a protozoan parasite that causes a neurological disease in horses called equine protozoal myeloencephalitis. The route of transmission is speculated to be by fecal-oral transfer of sporocysts shed from opossums. Controversy exists regarding both the natural life cycle for this parasite as well as the species identity of opossum *Sarcocystis*. To provide stage-specific material for species comparison, 27 opossums from southern Michigan were screened for *Sarcocystis* spp. sporocysts. Seven opossums were positive for *Sarcocystis* sporocysts by fecal flotation. A simplified, effective technique for isolation, excystation, and culture of opossum *Sarcocystis* sp. from mucosal scrapings was developed. All 7 *Sarcocystis* sp. isolates were successfully cultured to grow long term in equine dermal cells to the merozoite stage. Merozoites were observed between 5 and 15 days after inoculation. In conclusion, opossums shed *Sarcocystis* sp. sporocysts that may be manipulated to excyst and grow in vitro in equine dermal cell lines to the merozoite stage using the simplified technique described.

=> s sarcocystis neurona

L16 905 SARCOCYSTIS NEURONA

=> s l16 and (12 kd or 13 kd or 14 kd or 15 kd or 16 kd or 17 kd or 18 kd or 19 kd or 20 kd)

7 FILES SEARCHED...

L17 2 L16 AND (12 KD OR 13 KD OR 14 KD OR 15 KD OR 16 KD OR 17 KD OR 18 KD OR 19 KD OR 20 KD)

=> d bib ab 1-2

L17 ANSWER 1 OF 2 MEDLINE

AN 2000152631 MEDLINE

DN 20152631 PubMed ID: 10690772

TI Improvement of western blot test specificity for detecting equine serum antibodies to **Sarcocystis neurona**.

AU Rossano M G; Mansfield L S; Kaneene J B; Murphy A J; Brown C M; Schott H C 2nd; Fox J C

CS Animal Health Diagnostic Laboratory, The Population Medicine Center, Michigan State University, East Lansing 48824, USA.

SO JOURNAL OF VETERINARY DIAGNOSTIC INVESTIGATION, (2000 Jan) 12 (1) 28-32. Journal code: 9011490. ISSN: 1040-6387.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 200003

ED Entered STN: 20000330

Last Updated on STN: 20000330

Entered Medline: 20000321

AB Equine protozoal myeloencephalitis (EPM) is a neurological disease of horses and ponies caused by the apicomplexan protozoan parasite **Sarcocystis neurona**. The purposes of this study were to develop the most stringent criteria possible for a positive test result, to estimate the sensitivity and specificity of the EPM Western blot antibody test, and to assess the ability of bovine antibodies to *Sarcocystis cruzi* to act as a blocking agent to minimize false-positive results in the western blot test for *S. neurona*. **Sarcocystis neurona** merozoites harvested from equine dermal cell culture were heat denatured, and the proteins were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis in a 12-20% linear gradient gel. Separated proteins were electrophoretically transferred to polyvinylidene fluoride membranes and blocked in 1% bovine serum albumin

and 0.5% Tween-Tris-buffered saline. Serum samples from 6 horses with *S. neurona* infections (confirmed by culture from neural tissue) and 57 horses without infections (horses from the Eastern Hemisphere, where *S. neurona* does not exist) were tested by Western blot. Horses from both groups had reactivity to the 62-, 30-, 16-, 13-, 11-, 10.5-, and 10-kD bands. Testing was repeated with another step. Blots were treated with bovine *S. cruzi* antibodies prior to loading the equine samples. After this modification of the Western blot test, positive infection status was significantly associated with reactivity to the 30- and 16-kD bands ( $P < 0.001$ , Fisher's exact test). The *S. cruzi* antibody-blocked Western blot had a sample sensitivity of 100% and sample specificity of 98%. It is concluded that the specificity of the Western blot test is improved by blocking proteins not specific to *S. neurona* and using reactivity to the 30- and 16-kD bands as the criterion for a positive test.

L17 ANSWER 2 OF 2 WPIDS (C) 2003 THOMSON DERWENT  
 AN 1999-571872 [48] WPIDS  
 DNN N1999-421433 DNC C1999-166894  
 TI Biologically pure culture of equine *Neospora*, used as source of vaccines and diagnostic reagents.  
 DC B04 C06 C07 D16 S03  
 IN BARR, B C; CONRAD, P A; MARSH, A E  
 PA (REGC) UNIV CALIFORNIA  
 CYC 23  
 PI WO 9947927 A1 19990923 (199948)\* EN 47p  
 RW: AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL PT SE  
 W: AU CA JP  
 AU 9931874 A 19991011 (200008)  
 US 6071737 A 20000606 (200033)  
 EP 1064550 A1 20010103 (200102) EN  
 R: AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE  
 JP 2002509702 W 20020402 (200225) 47p  
 ADT WO 9947927 A1 WO 1999-US5754 19990316; AU 9931874 A AU 1999-31874 19990316; US 6071737 A US 1998-42600 19980316; EP 1064550 A1 EP 1999-913906 19990316, WO 1999-US5754 19990316; JP 2002509702 W WO 1999-US5754 19990316, JP 2000-537071 19990316  
 FDT AU 9931874 A Based on WO 9947927; EP 1064550 A1 Based on WO 9947927; JP 2002509702 W Based on WO 9947927  
 PRAI US 1998-42600 19980316  
 AB WO 9947927 A UPAB: 19991122  
 NOVELTY - Biologically pure culture of equine *Neospora*, is new.  
 DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following:  
 (a) detecting antibodies (Ab) specifically reactive with equine *Neospora* antigens (Ag) by forming an Ab-Ag complex;  
 (b) detecting *Neospora* by forming a complex with an antibody (Ab1) specifically reactive with *Neospora* antigen;  
 (c) detecting *Neospora*-specific nucleic acid (I) by hybridization with a specific oligonucleotide probe; and  
 (d) pharmaceutical composition containing equine *Neospora* immunogen and a carrier.  
 ACTIVITY - Antiprotozoal.  
 MECHANISM OF ACTION - Induction of a specific immune response.  
 USE - Immunogens (optionally expressed from gene therapy vectors) from equine *Neospora* are used in vaccines for treatment or prevention of *Neospora* infection in horses and other animals. *Neospora* is a causative agent of equine protozoal myeloencephalitis (EPM). Detection of *Neospora*-specific antigens, antibodies or nucleic acid (by usual immunoassay or hybridization tests) is used to diagnose infection. Antibodies (Ab) specific for equine *Neospora* are used for diagnosis; to select candidate immunogens for vaccine development; to isolate proteins; to screen DNA libraries and as therapeutic/prophylactic agents.

ADVANTAGE - Reagents specific for equine Neospora allow differentiation between equine protozoal myeloencephalitis caused by Neospora and **Sarcocystis neurona**. These pathogens require different treatments and treatment of Neospora is only effective if applied before the parasite has formed cysts. The vaccines also prevent shedding of oocysts by animals known to be infected.  
Dwg.0/2

=> s l16 and (26 kd or 27 kd or 28 kd or 29 kd or 30 kd or 31 kd or 32 kd or 33 kd or 34 kd)

L18            0 L16 AND (26 KD OR 27 KD OR 28 KD OR 29 KD OR 30 KD OR 31 KD OR 32 KD OR 33 KD OR 34 KD)

=> d his

(FILE 'HOME' ENTERED AT 14:49:07 ON 15 MAY 2003)

FILE 'BIOSIS, MEDLINE, AGRICOLA, EMBASE, CABA, WPIDS, JAPIO, BIOTECHDS, LIFESCI, CAPLUS' ENTERED AT 14:49:30 ON 15 MAY 2003

                  E MANSFIELD LINDA S/AU  
L1                30 S E2-E3  
                  E MANSFIELD L S/AU  
L2                139 S E3  
                  E ROSSANO MARY G/AU  
L3                9 S E3  
                  E ROSSANO M G/AU  
L4                25 S E3  
                  E MURPHY ALICE J/AU  
L5                15 S E2 OR E3  
                  E MURPHY A J/AU  
L6                337 S E3  
                  E VRABLE RUTH A/AU  
L7                11 S E2 OR E3  
                  E VRABLE R A/AU  
L8                23 S E2 OR E3  
L9                496 S L1-L8  
L10               39 S L9 AND SARCOCYSTIS NEURONA  
L11               1 S L10 AND (16 KD OR 30 KD)  
L12               51 S L9 AND ANTIBOD?  
L13               26 S L10 AND ANTIBOD?  
L14               12 DUP REM L13 (14 DUPLICATES REMOVED)  
L15               15 DUP REM L10 (24 DUPLICATES REMOVED)  
L16               905 S SARCOCYSTIS NEURONA  
L17               2 S L16 AND (12 KD OR 13 KD OR 14 KD OR 15 KD OR 16 KD OR 17 KD  
L18               0 S L16 AND (26 KD OR 27 KD OR 28 KD OR 29 KD OR 30 KD OR 31 KD

=> s l16 and molecular weight

L19               5 L16 AND MOLECULAR WEIGHT

=> s l19 and antibod?

L20               3 L19 AND ANTIBOD?

=> d bib ab 1-3

L20 ANSWER 1 OF 3            MEDLINE  
AN 2002085433            MEDLINE  
DN 21671299    PubMed ID: 11812499  
TI Molecular characterisation of a major 29 kDa surface antigen of  
**Sarcocystis neurona**.  
AU Ellison Siobhan P; Omara-Opyene A Levi; Yowell Charles A; Marsh Antoinette  
E; Dame John B  
CS Department of Pathobiology, University of Florida, P.O. Box 110880,

Gainesville, FL 32611-0880, USA.

SO INTERNATIONAL JOURNAL FOR PARASITOLOGY, (2002 Feb) 32 (2) 217-25.  
Journal code: 0314024. ISSN: 0020-7519.

CY England; United Kingdom

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

OS GENBANK-AF397896; GENBANK-AF401682

EM 200205

ED Entered STN: 20020129  
Last Updated on STN: 20020517  
Entered Medline: 20020516

AB A gene encoding a major 29 kDa surface antigen from **Sarcocystis neurona**, the primary causative agent of equine protozoal myeloencephalitis (EPM), was cloned, sequenced, and expressed as a recombinant protein. A cDNA library was prepared in the expression vector lambda ZAP from polyA+mRNA isolated from *S. neurona* merozoites cultivated in vitro. Random sequencing of 96 clones identified a clone of an abundant transcript having a translated amino acid sequence with 30% identity to the 31-kDa surface antigen of *Sarcocystis muris* cyst merozoites. Southern blot analysis indicated that the corresponding gene exists in low copy number within the *S. neurona* genome, but RNA blot analysis and other data indicated that the gene transcript is highly abundant. The sequence of the cDNA clone encoded an open reading frame specifying a polypeptide of 276 amino acids with a predicted size of 28.7 kDa. The deduced amino acid sequence displayed a hypothetical N-terminal signal peptide sequence followed by a polypeptide containing 12 cysteines. The coding region of the cDNA insert was subcloned into the expression vector pET14b, and a fusion protein expressed. The recombinant polypeptide was recognised by mAb 2A7 and mAb 1631, directed against a 29 kDa native protein found on the surface of cultured merozoites. **Antibodies** in serum and cerebrospinal fluid from a horse with EPM recognised a 29 kDa native protein of *S. neurona* merozoites and the 29 kDa recombinant protein. This *S. neurona* surface antigen is named SnSAG1.

L20 ANSWER 2 OF 3 MEDLINE

AN 2001354018 MEDLINE

DN 21127318 PubMed ID: 11223200

TI Immunoconversion against **Sarcocystis neurona** in normal and dexamethasone-treated horses challenged with *S. neurona* sporocysts.

AU Cutler T J; MacKay R J; Ginn P E; Gillis K; Tanhauser S M; LeRay E V; Dame J B; Greiner E C

CS Department of Pathobiology, PO Box 100880, College of Veterinary Medicine, University of Florida, Gainesville 32610, USA.

SO VETERINARY PARASITOLOGY, (2001 Feb 26) 95 (2-4) 197-210.  
Journal code: 7602745. ISSN: 0304-4017.

CY Netherlands

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 200106

ED Entered STN: 20010625  
Last Updated on STN: 20010625  
Entered Medline: 20010621

AB Equine protozoal myeloencephalitis is a common neurologic disease of horses in the Americas usually caused by **Sarcocystis neurona**. To date, the disease has not been induced in horses using characterized sporocysts from *Didelphis virginiana*, the definitive host. *S. neurona* sporocysts from 15 naturally infected opossums were fed to horses seronegative for **antibodies** against *S. neurona*. Eight horses were given 5x10(5) sporocysts daily for 7 days. Horses were examined for abnormal clinical signs, and blood and cerebrospinal fluid were harvested at intervals for 90 days after the first day of challenge

=> s 116 and kd  
L22 3 L16 AND KD

=> d bib ab 1-3

L22 ANSWER 1 OF 3 MEDLINE  
AN 2000152631 MEDLINE  
DN 20152631 PubMed ID: 10690772  
TI Improvement of western blot test specificity for detecting equine serum antibodies to **Sarcocystis neurona**.  
AU Rossano M G; Mansfield L S; Kaneene J B; Murphy A J; Brown C M; Schott H C 2nd; Fox J C  
CS Animal Health Diagnostic Laboratory, The Population Medicine Center, Michigan State University, East Lansing 48824, USA.  
SO JOURNAL OF VETERINARY DIAGNOSTIC INVESTIGATION, (2000 Jan) 12 (1) 28-32. Journal code: 9011490. ISSN: 1040-6387.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200003  
ED Entered STN: 20000330  
Last Updated on STN: 20000330  
Entered Medline: 20000321  
AB Equine protozoal myeloencephalitis (EPM) is a neurological disease of horses and ponies caused by the apicomplexan protozoan parasite **Sarcocystis neurona**. The purposes of this study were to develop the most stringent criteria possible for a positive test result, to estimate the sensitivity and specificity of the EPM Western blot antibody test, and to assess the ability of bovine antibodies to **Sarcocystis cruzi** to act as a blocking agent to minimize false-positive results in the western blot test for **S. neurona**. **Sarcocystis neurona** merozoites harvested from equine dermal cell culture were heat denatured, and the proteins were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis in a 12-20% linear gradient gel. Separated proteins were electrophoretically transferred to polyvinylidene fluoride membranes and blocked in 1% bovine serum albumin and 0.5% Tween-Tris-buffered saline. Serum samples from 6 horses with **S. neurona** infections (confirmed by culture from neural tissue) and 57 horses without infections (horses from the Eastern Hemisphere, where **S. neurona** does not exist) were tested by Western blot. Horses from both groups had reactivity to the 62-, 30-, 16-, 13-, 11-, 10.5-, and 10-kD bands. Testing was repeated with another step. Blots were treated with bovine **S. cruzi** antibodies prior to loading the equine samples. After this modification of the Western blot test, positive infection status was significantly associated with reactivity to the 30- and 16-kD bands ( $P < 0.001$ , Fisher's exact test). The **S. cruzi** antibody-blocked Western blot had a sample sensitivity of 100% and sample specificity of 98%. It is concluded that the specificity of the Western blot test is improved by blocking proteins not specific to **S. neurona** and using reactivity to the 30- and 16-kD bands as the criterion for a positive test.

L22 ANSWER 2 OF 3 MEDLINE  
AN 93222344 MEDLINE  
DN 93222344 PubMed ID: 8466988  
TI Equine protozoal myeloencephalitis: antigen analysis of cultured **Sarcocystis neurona** merozoites.  
AU Granstrom D E; Dubey J P; Davis S W; Fayer R; Fox J C; Poonacha K B; Giles R C; Comer P F  
CS Department of Veterinary Science, University of Kentucky, Lexington 40546-0099.

SO JOURNAL OF VETERINARY DIAGNOSTIC INVESTIGATION, (1993 Jan) 5 (1) 88-90.  
 Journal code: 9011490. ISSN: 1040-6387.

CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199305  
 ED Entered STN: 19930521  
 Last Updated on STN: 19930521  
 Entered Medline: 19930510

AB Antigens of cultured *Sarcocystis neurona* merozoites were examined using immunoblot analysis. Blotted proteins were probed with *S. cruzi*, *S. muris*, and *S. neurona* antisera produced in rabbits, *S. fayeri* (pre- and post-infection) and *S. neurona* (pre- and post-inoculation) sera produced in horses, immune sera from 7 histologically confirmed cases of equine protozoal myeloencephalitis (EPM), and pre-suckle serum from a newborn foal. Eight proteins, 70, 24, 23.5, 22.5, 13, 11, 10.5, and 10 Kd, were detected only by *S. neurona* antiserum and/or immune serum from EPM-affected horses. Equine sera were titrated by the indirect immunofluorescent antibody (IFA) method using air-dried, cultured *S. neurona* merozoites. Anti-*Sarcocystis* IFA titers were found in horses with or without EPM. Serum titers did not correspond to the number of specific bands recognized on immunoblots.

L22 ANSWER 3 OF 3 WPIDS (C) 2003 THOMSON DERWENT  
 AN 1999-571872 [48] WPIDS  
 DNN N1999-421433 DNC C1999-166894  
 TI Biologically pure culture of equine Neospora, used as source of vaccines and diagnostic reagents.  
 DC B04 C06 C07 D16 S03  
 IN BARR, B C; CONRAD, P A; MARSH, A E  
 PA (REGC) UNIV CALIFORNIA  
 CYC 23  
 PI WO 9947927 A1 19990923 (199948)\* EN 47p  
 RW: AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL PT SE  
 W: AU CA JP  
 AU 9931874 A 19991011 (200008)  
 US 6071737 A 20000606 (200033)  
 EP 1064550 A1 20010103 (200102) EN  
 R: AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE  
 JP 2002509702 W 20020402 (200225) 47p

ADT WO 9947927 A1 WO 1999-US5754 19990316; AU 9931874 A AU 1999-31874 19990316; US 6071737 A US 1998-42600 19980316; EP 1064550 A1 EP 1999-913906 19990316, WO 1999-US5754 19990316; JP 2002509702 W WO 1999-US5754 19990316, JP 2000-537071 19990316

FDT AU 9931874 A Based on WO 9947927; EP 1064550 A1 Based on WO 9947927; JP 2002509702 W Based on WO 9947927

PRAI US 1998-42600 19980316

AB WO 9947927 A UPAB: 19991122  
 NOVELTY - Biologically pure culture of equine Neospora, is new.  
 DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following:  
 (a) detecting antibodies (Ab) specifically reactive with equine Neospora antigens (Ag) by forming an Ab-Ag complex;  
 (b) detecting Neospora by forming a complex with an antibody (Ab1) specifically reactive with Neospora antigen;  
 (c) detecting Neospora-specific nucleic acid (I) by hybridization with a specific oligonucleotide probe; and  
 (d) pharmaceutical composition containing equine Neospora immunogen and a carrier.  
 ACTIVITY - Antiprotozoal.  
 MECHANISM OF ACTION - Induction of a specific immune response.  
 USE - Immunogens (optionally expressed from gene therapy vectors)

from equine Neospora are used in vaccines for treatment or prevention of Neospora infection in horses and other animals. Neospora is a causative agent of equine protozoal myeloencephalitis (EPM). Detection of Neospora-specific antigens, antibodies or nucleic acid (by usual immunoassay or hybridization tests) is used to diagnose infection. Antibodies (Ab) specific for equine Neospora are used for diagnosis; to select candidate immunogens for vaccine development; to isolate proteins; to screen DNA libraries and as therapeutic/prophylactic agents.

ADVANTAGE - Reagents specific for equine Neospora allow differentiation between equine protozoal myeloencephalitis caused by Neospora and **Sarcocystis neurona**. These pathogens require different treatments and treatment of Neospora is only effective if applied before the parasite has formed cysts. The vaccines also prevent shedding of oocysts by animals known to be infected.

Dwg.0/2

=> s l16 and antibod?

L23 343 L16 AND ANTIBOD?

=> s l23 and (treat? or therap?)

4 FILES SEARCHED...

L24 50 L23 AND (TREAT? OR THERAP?)

=> dup rem l24

PROCESSING COMPLETED FOR L24

L25 23 DUP REM L24 (27 DUPLICATES REMOVED)

=> d bib ab 1-23

L25 ANSWER 1 OF 23 CAPLUS COPYRIGHT 2003 ACS

AN 2002:946917 CAPLUS

DN 138:3686

TI Monoclonal **antibodies** to **Sarcocystis neurona**  
and uses thereof

IN Marsh, Antoinette

PA USA

SO U.S. Pat. Appl. Publ., 14 pp., which

CODEN: USXXCO

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2002187517	A1	20021212	US 2002-140754	20020507
PRAI	US 2001-293603P	P	20010524		
	US 2001-297810P	P	20010612		

AB The present invention is directed to particular monoclonal **antibodies** (2A7-18 and 2G5-2) that find use in the identification and purifn. of *S. neurona* and related antigens. In particular, these **antibodies** permit the diagnosis of *Sarcocystis* related diseases such as equine protozoal myeloencephalitis (EPM).

L25 ANSWER 2 OF 23 CAPLUS COPYRIGHT 2003 ACS

AN 2002:638328 CAPLUS

DN 137:151587

TI Use of SAG-1 gene of **Sarcocystis neurona** for  
diagnostic tests and vaccines for equine protozoal myeloencephalitis

IN Dame, John B.; Ellison, Siobhan P.; Yowell, Charles A.

PA USA

SO U.S. Pat. Appl. Publ., 21 pp.

CODEN: USXXCO

DT Patent



LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2002115828	A1	20020822	US 2001-962993	20010924
PRAI	US 2000-234676P	P	20000922		

AB A gene encoding a 29 kilodalton protein found on the surface of merozoite stage *S. neurona* has been cloned and sequenced. The protein encoded by this gene, termed SnSAG-1, is an immunodominant antigen recognized on protein blots. Methods for using nucleic acids and polypeptides relating to SnSAG-1 in diagnostic tests and vaccine development are disclosed. Claimed sequences were not present at the time of publication.

L25 ANSWER 3 OF 23 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE 1

AN 2003:112305 BIOSIS

DN PREV200300112305

TI Experimental induction of equine protozoan myeloencephalitis (EPM) in the horse: Effect of *Sarcocystis neurona* sporocyst inoculation dose on the development of clinical neurologic disease.

AU Sofaly, C. D. (1); Reed, S. M.; Gordon, J. C. (1); Dubey, J. P.; Oglesbee, M. J.; Njoku, C. J. (1); Grover, D. L. (1); Saville, W. J. A. (1)

CS (1) Department of Veterinary Preventive Medicine, College of Veterinary Medicine, The Ohio State University, Columbus, OH, 43210-1092, USA: saville.4@osu.edu USA

SO Journal of Parasitology, (December 2002, 2002) Vol. 88, No. 6, pp. 1164-1170. print. ISSN: 0022-3395.

DT Article

LA English

AB The effect of inoculation dose of *Sarcocystis neurona* sporocysts on the development of clinical neurologic disease in horses was investigated. Twenty-four seronegative weanling horses were subjected to the natural stress of transport and then randomly assigned to 6 treatment groups of 4 horses each. Horses were then immediately inoculated with either 102, 103, 104, 105, or 106 *S. neurona* sporocysts or placebo using nasogastric tube and housed indoors. Weekly neurologic examinations were performed by a blinded observer. Blood was collected weekly for antibody determination by Western blot analysis. Cerebrospinal fluid was collected before inoculation and before euthanasia for *S. neurona* antibody determination. Horses were killed and necropsied between 4 and 5 wk after inoculation. Differences were detected among dose groups based on seroconversion times, severity of clinical neurologic signs, and presence of microscopic lesions. Seroconversion of challenged horses was observed as early as 14 days postinfection in the 106 sporocyst dose group. Mild to moderate clinical signs of neurologic disease were produced in challenged horses from all groups, with the most consistent signs seen in the 106 sporocyst dose group. Histologic lesions suggestive of *S. neurona* infection were detected in 4 of the 20 horses fed sporocysts. Parasites were not detected in equine tissues by light microscopy, immunohistochemistry, or bioassay in gamma-interferon gene knockout mice. Control horses remained seronegative for the duration of the study and had no histologic evidence of protozoal infection.

L25 ANSWER 4 OF 23 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE 2

AN 2002:348434 BIOSIS

DN PREV200200348434

TI Reduced levels of nitric oxide metabolites in cerebrospinal fluid are associated with equine protozoal myeloencephalitis.

AU Njoku, Chinedu J.; Saville, William J. A.; Reed, Stephen M.; Oglesbee, Michael J.; Rajala-Schultz, Paivi J.; Stich, Roger W. (1)

CS (1) Department of Veterinary Preventive Medicine, The Ohio State

University, 1900 Coffey Rd., Columbus, OH, 43210-1092: stich.2@osu.edu USA  
 SO Clinical and Diagnostic Laboratory Immunology, (May, 2002) Vol. 9, No. 3,  
 pp. 605-610. print.  
 ISSN: 1071-412X.  
 DT Article  
 LA English  
 AB Equine protozoal myeloencephalitis (EPM) is a disease of horses that is  
 primarily associated with infection with the apicomplexan  
**Sarcocystis neurona**. Infection with this parasite alone  
 is not sufficient to induce the disease, and the mechanism of  
 neuropathogenesis associated with EPM has not been reported. Nitric oxide  
 (NO) functions as a neurotransmitter, a vasodilator, and an immune  
 effector and is produced in response to several parasitic protozoa. The  
 purpose of this work was to determine if the concentration of NO  
 metabolites (NOx-) in the cerebrospinal fluid (CSF) is correlated with the  
 development of EPM. CSF NOx- levels were measured before and after  
 transport-stressed, acclimated, or dexamethasone-**treated** horses  
 (n = 3 per group) were experimentally infected with *S. neurona* sporocysts.  
 CSF NOx- levels were also compared between horses that were diagnosed with  
 EPM after natural infection with *S. neurona* and horses that did not have  
 clinical signs of disease or that showed no evidence of infection with the  
 parasite (n = 105). Among the experimentally infected animals, the mean  
 CSF NOx- levels of the transport-stressed group, which had the most severe  
 clinical signs, was reduced after infection, while these values were found  
 to increase after infection in the remaining groups that had less severe  
 signs of EPM. Under natural conditions, horses with EPM (n = 65) had a  
 lower mean CSF NOx- concentration than clinically normal horses with  
**antibodies** (Abs) against *S. neurona* (n = 15) in CSF, and horses  
 that developed ataxia (n = 81) had a significantly lower mean CSF NOx-  
 concentration than horses that did not have neurologic signs (n = 24). In  
 conclusion, lower CSF NOx- levels were associated with clinical EPM,  
 suggesting that measurement of CSF NOx- levels could improve the accuracy  
 of diagnostic tests that are based upon detection of *S. neurona*-specific  
 Abs in CSF alone and that reduced NO levels could be causatively related  
 to the development of EPM.

L25 ANSWER 5 OF 23 CABA COPYRIGHT 2003 CABI  
 AN 2002:114296 CABA  
 DN 20023070361  
 TI Folate deficiency during **treatment** with orally administered  
 folic acid, sulphadiazine and pyrimethamine in a horse with suspected  
 equine protozoal myeloencephalitis (EPM)  
 AU Piercy, R. J.; Hinchcliff, K. W.; Reed, S. M.  
 CS Department of Clinical Veterinary Science, The Ohio State University, 601  
 Vernon L. Tharp Street, Columbus, OH 43210, USA.  
 SO Equine Veterinary Journal, (2002) Vol. 34, No. 3, pp. 311-316. 35 ref.  
 ISSN: 0425-1644  
 DT Journal  
 LA English  
 AB A 6-year-old Quarter Horse show-mare with tail block suffering from  
 dysphagia and ataxia believed to be the cause of equine protozoal  
 myeloencephalitis (EPM) is presented at the Department of Clinical  
 Veterinary Science, Ohio State University, Columbus Ohio, USA [date not  
 given]. Sulfadiazine (14.7-44.4 mg/kg), pyrimethamine (0.7-2.2 mg/kg) and  
 folic acid (9.6 mg, q. 12 h) was given to the animal for 9 months.  
 Physical examination revealed ulceration and inflammation of the tongue,  
 and a draining abscess and localized subcutaneous swelling at the dorsal  
 aspect of the tail head. Haematology and serum biochemistry revealed  
 anaemia and leukopenia, neutropenia, lymphopenia, low total CO<sub>2</sub>,  
 hyperbilirubinaemia, and elevated creatinine and albumin. Previous  
 medications were stopped and lactated Ringer's solution (120 ml/kg/day,  
 i.v.), potassium penicillin (22 000 IU/kg, q 6 h, i.v.), gentamicin  
 (Gentocin, 6.6 mg/kg, i.v.) and folic acid (0.11 mg/kg, i.v.) were

administered to the horse. After 6 days of **treatment**, cerebrospinal fluid and serum were submitted for **Sarcocystis neurona antibodies**. On day 8, there was complete healing of lingual ulcers and penicillin and gentamicin were discontinued while ceftiofur (Naxcel, 2.2 mg/kg q 12 h, i.m.) was administered. Four days later, the presence of **S. neurona antibodies** was confirmed, thus, supporting the diagnosis of EPM, and diclazuril (Clinacox, 5 mg/kg q 24 h, p.o.) was administered. After 68 days, the physical, oral and neurological examinations were declared all normal. It is concluded that folic acid supplementation does not prevent the development of folate deficiency and that it is not recommended to supplement folic acid to horses being **treated** with dihydrofolate reductase inhibitors.

L25 ANSWER 6 OF 23 WPIDS (C) 2003 THOMSON DERWENT  
AN 2002-049244 [06] WPIDS  
DNC C2002-013806  
TI Vaccine useful for preventing or ameliorating equine protozoal myeloencephalitis disease, comprises inactivated **Sarcocystis neurona** cells and/or Neospora hughesi cells, antigens, DNA derived from the cells or their mixtures.  
DC B04 C06 D16  
IN BIGBIE, R B; NG, T K; WHALEN, J W  
PA (AMHP) AMERICAN HOME PROD CORP; (AMHP) WYETH  
CYC 96  
PI WO 2001080885 A2 20011101 (200206)\* EN 31p  
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ  
NL OA PT SD SE SL SZ TR TZ UG ZW  
W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK  
DM DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ  
LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD  
SE SG SI SK SL TJ TM TR TT TZ UA UG UZ VN YU ZA ZW  
AU 2001051761 A 20011107 (200219)  
US 2002041886 A1 20020411 (200227)  
EP 1276499 A2 20030122 (200308) EN  
R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
RO SE SI TR  
BR 2001010232 A 20030121 (200309)  
ADT WO 2001080885 A2 WO 2001-US40527 20010413; AU 2001051761 A AU 2001-51761  
20010413; US 2002041886 A1 Provisional US 2000-199435P 20000425,  
Provisional US 2001-278695P 20010326, US 2001-840485 20010423; EP 1276499  
A2 EP 2001-925175 20010413, WO 2001-US40527 20010413; BR 2001010232 A BR  
2001-10232 20010413, WO 2001-US40527 20010413  
FDT AU 2001051761 A Based on WO 2001080885; EP 1276499 A2 Based on WO  
2001080885; BR 2001010232 A Based on WO 2001080885  
PRAI US 2001-278695P 20010326; US 2000-199435P 20000425; US 2001-840485  
20010423  
AB WO 2001080885 A UPAB: 20020128  
NOVELTY - An immunogenically active component (I) comprising a merozoite  
**antibody** inducing, inactivated **Sarcocystis**  
**neurona** cells, tachyzoite **antibody** inducing, inactivated  
Neospora hughesi cells, a merozoite or tachyzoite **antibody**  
inducing antigen derived from the cells, DNA derived from the cells  
capable of inducing a merozoite or tachyzoite **antibody** immune  
response or their mixture, is new.  
DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the  
following:  
(1) a vaccine composition comprising (I), a pharmacologically  
acceptable carrier, and an immunogenically stimulating adjuvant;  
(2) cell culture propagation (III) of *S. neurona* or *N. hughesi*  
protozoan parasite, comprising:  
(i) growing a monolayer of cells having a confluency of 80-100%;  
(ii) re-feeding the cells with supplemented growth media;  
(iii) inoculating the cells with merozoites or tachyzoites;

(iv) holding the inoculated cells for 4-12 days; and  
(v) decanting the supplemented growth media from the inoculated cells and refeeding the cells a second time with supplemented growth media; and  
(3) preventing or ameliorating an EPM disease in equines, comprising administering an immunogenically active component selected from the following:

- (i) merozoite **antibody** inducing, inactivated **Sarcocystis neurona** cells;
- (ii) tachyzoite **antibody** inducing, inactivated *Neospora hughesi* cells;
- (iii) a merozoite or tachyzoite **antibody** inducing antigen derived from the cells;
- (iv) DNA derived from the cells, capable of inducing a merozoite or tachyzoite **antibody** immune response; or
- (v) a mixture of the above.

ACTIVITY - Neuroprotective; Antiinflammatory. No biological data was provided.

MECHANISM OF ACTION - Vaccine. One group of horses were administered 1 multiply 105 merozoites/dose of the vaccine, a second group of 21 horses were administered vaccine blended at 1 multiply 106 merozoites/dose, a third group of 10 horses were administered vaccine at 1 multiply 107 merozoites/dose, and a fourth of group of 10 horses were maintained as non-vaccinated environmental controls. **Treated** horses from all groups showed significant increases in **antibodies** to *S. neurona* merozoites while the control horses maintained a low to non-existent **antibody** level.

USE - (I) and (II) are useful for prevention or amelioration of equine protozoal myeloencephalitis (EPM) disease in equines. (III) is useful for the propagation of cells such as equine dermal, maiden darby bovine kidney, African green monkey kidney, canine monocyte, mouse monocyte, fetal rhesus monkey kidney, feline kidney, maiden darby canine kidney and baby hamster kidney cells (claimed).  
Dwg:0/0

L25 ANSWER 7 OF 23 BIOTECHDS COPYRIGHT 2003 THOMSON DERWENT AND ISI  
AN 2002-04497 BIOTECHDS  
TI Vaccine useful for preventing or ameliorating equine protozoal myeloencephalitis disease, comprises inactivated *Sarcocystis neurona* cells and/or *Neospora hughesi* cells, antigens, DNA derived from the cells or their mixtures;  
horse protozoan myeloencephalitis disease **therapy** using a recombinant vaccine or a nucleic acid vaccine  
AU Bigbie R B; Ng T K; Whalen Jr J W  
PA American-Home-Prod.  
LO Madison, NJ, USA.  
PI WO 2001080885 1 Nov 2001  
AI WO 2001-US40527 13 Apr 2001  
PRAI US 2001-278695 26 Mar 2001; US 2000-199435 25 Apr 2000  
DT Patent  
LA English  
OS WPI: 2002-049244 [06]  
AB An immunogenically active component (I) having a merozoite **antibody** inducing, inactivated **Sarcocystis neurona** cells, tachyzoite **antibody** inducing, inactivated *Neospora hughesi* cells, a merozoite or tachyzoite **antibody** inducing antigen derived from the cells, DNA derived from the cells capable of inducing a merozoite or tachyzoite **antibody** immune response or their mixture, is new. Also claimed are: a vaccine composition comprising (I), a pharmacologically acceptable carrier, and an immunogenically stimulating adjuvant; cell culture propagation (III) of *S. neurona* or *N. hughesi* protozoan parasite, involving growing a monolayer of cells, re-feeding with supplemented growth medium, inoculating with merozoites or tachyzoites, holding for

4-12 days and decanting the supplemented growth medium from the inoculated cells and refeeding a second time with supplemented growth medium; and preventing or ameliorating an EPM disease in horses. (III) is useful for the propagation of cells such as horse dermal, cattle kidney, African green monkey kidney, dog monocyte, mouse monocyte, fetal rhesus monkey kidney, feline kidney, dog kidney and baby hamster kidney cells. (31pp)

L25 ANSWER 8 OF 23 EMBASE COPYRIGHT 2003 ELSEVIER SCI. B.V.  
 AN 2001075088 EMBASE  
 TI Migration and development of **Sarcocystis neurona** in tissues of interferon gamma knockout mice fed sporocysts from a naturally infected opossum.  
 AU Dubey J.P.  
 CS J.P. Dubey, United States Dept. of Agriculture, Parasite Biol., Epidem./Syst. Lab., Animal and Natural Resources Inst, Beltsville, MD 20705-2350, United States. jdubey@anri.barc.usda.gov  
 SO Veterinary Parasitology, (26 Feb 2001) 95/2-4 (341-351).  
 Refs: 19  
 ISSN: 0304-4017 CODEN: VPARDI  
 PUI S 0304-4017(00)00401-5  
 CY Netherlands  
 DT Journal; Article  
 FS 004 Microbiology  
 005 General Pathology and Pathological Anatomy  
 LA English  
 SL English  
 AB Migration and development of **Sarcocystis neurona** was studied in 50 gamma interferon knockout mice fed graded doses of S. neurona sporocysts from the intestine of a naturally infected opossum. Mice were examined at necropsy 1-62 days after feeding sporocysts (DAFS). All tissue sections were reacted with anti-S. neurona-specific polyclonal rabbit serum in an immunohistochemical (IHC) test. Between 1 and 3 DAFS, organisms were seen mainly in intestines. Between 4 and 11 DAFS, organisms were seen in several visceral tissues. Beginning with 13 DAFS, schizonts and merozoites were present in sections of brains of all infected mice. All regions of the brain were parasitized but the hind brain was most severely affected. S. neurona was found in the spinal cord of all 10 mice examined 22-30 DAFS. Of the 28 infected mice examined 20-62 DAFS, S. neurona was found in the brains of all 28, lungs of 14, hearts of 8 and eyes of 3. More organisms were seen in IHC-stained sections than in sections stained with hematoxylin and eosin. **Treatment** of tissues with glutaraldehyde, Karnovsky fixative, and ethylene diamino tetra acetic acid (EDTA, used for decalcification) did not affect staining of organisms by IHC.

L25 ANSWER 9 OF 23 MEDLINE DUPLICATE 3  
 AN 2001511543 MEDLINE  
 DN 21442379 PubMed ID: 11558662  
 TI Suspected protozoal myeloencephalitis in a two-month-old colt.  
 AU Gray L C; Magdesian K G; Sturges B K; Madigan J E  
 CS Veterinary Medical Teaching Hospital, Large Animal Clinic, Department of Medicine and Epidemiology, School of Veterinary Medicine, University of California, Davis 95616, USA.  
 SO VETERINARY RECORD, (2001 Sep 1) 149 (9) 269-73.  
 Journal code: 0031164. ISSN: 0042-4900.  
 CY England: United Kingdom  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 200201  
 ED Entered STN: 20010918  
 Last Updated on STN: 20020125

horse would have clinical improvement. The likelihood of survival among horses with EPM was lower among horses with more severe clinical signs and higher among horses that improved after EPM was diagnosed. Conclusions and Clinical Relevance: **Treatment** of horses with EPM is indicated in most situations; however, severity of clinical signs should be taken into consideration when making **treatment** decisions. Response to **treatment** is an important indicator of survival.

L25 ANSWER 14 OF 23 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE 7

AN 2000:339710 BIOSIS

DN PREV200000339710

TI Detection of **Sarcocystis neurona** in the brain of a Grant's zebra (*Equus burchelli bohmi*).

AU Marsh, Antoinette E. (1); Denver, Mary; Hill, Frazer I.; McElhaney, M. R.; Trupkiewicz, J. G.; Stewart, James; Tell, Lisa

CS (1) Department of Veterinary Pathobiology, University of Missouri, Columbia, MO, 65211 USA

SO Journal of Zoo and Wildlife Medicine, (March, 2000) Vol. 31, No. 1, pp. 82-86. print.  
ISSN: 1042-7260.

DT Article

LA English

SL English

AB An 8-yr-old intact male Grant's zebra (*Equus burchelli bohmi*) was referred to the Veterinary Medical Teaching Hospital of the University of California-Davis after being found in the owner's pasture obtunded and in lateral recumbency. The animal was hypothermic, weak, and unwilling to rise. There was no evidence of trauma, and the zebra had seemed normal the preceding evening. There was no extensor rigidity, and cranial nerve reflexes were normal. Flexor and extensor reflexes were weak upon initial examination. A complete blood count and serum biochemistry analysis revealed a mild leukocytosis, hyperfibrinogenemia, hypoglycemia, hyponatremia, hypochloremia, hypocalcemia, and hypoalbuminemia. Urinalysis was normal, and a urine toxicology screen for alkaloids was negative. No toxic substance was found in the hay or pasture grasses although the owner reported the presence of yellow star thistle and mushrooms in the pasture. The cerebrospinal fluid cytologic and biochemical analyses were normal, but **antibodies to Sarcocystis neurona** were detected. The zebra died despite aggressive supportive **therapy** over a 4-day period. The necropsy demonstrated severe gastrointestinal nematodiasis that could account for hypoalbuminemia and electrolyte abnormalities. Histopathologic examination of the nervous system revealed focal areas of perivascular cuffing in the brainstem that were comprised mainly of lymphocytes, monocytes, and plasma cells. Immunohistochemical staining identified the presence of *S. neurona* merozoites associated with the lesions. This zebra probably died from severe endoparasitism that resulted in malabsorption, weakness, and recumbency rather than from encephalitis associated with *S. neurona* merozoites. Equine protozoal myeloencephalitis has not been reported previously in nondomestic equids.

L25 ANSWER 15 OF 23 MEDLINE DUPLICATE 8

AN 2000152631 MEDLINE

DN 20152631 PubMed ID: 10690772

TI Improvement of western blot test specificity for detecting equine serum **antibodies to Sarcocystis neurona**.

AU Rossano M G; Mansfield L S; Kaneene J B; Murphy A J; Brown C M; Schott H C 2nd; Fox J C

CS Animal Health Diagnostic Laboratory, The Population Medicine Center, Michigan State University, East Lansing 48824, USA.

SO JOURNAL OF VETERINARY DIAGNOSTIC INVESTIGATION, (2000 Jan) 12 (1) 28-32.  
Journal code: 9011490. ISSN: 1040-6387.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200003  
ED Entered STN: 20000330

Last Updated on STN: 20000330  
Entered Medline: 20000321

AB Equine protozoal myeloencephalitis (EPM) is a neurological disease of horses and ponies caused by the apicomplexan protozoan parasite **Sarcocystis neurona**. The purposes of this study were to develop the most stringent criteria possible for a positive test result, to estimate the sensitivity and specificity of the EPM Western blot **antibody** test, and to assess the ability of bovine **antibodies** to **Sarcocystis cruzi** to act as a blocking agent to minimize false-positive results in the western blot test for **S. neurona**. **Sarcocystis neurona** merozoites harvested from equine dermal cell culture were heat denatured, and the proteins were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis in a 12-20% linear gradient gel. Separated proteins were electrophoretically transferred to polyvinylidene fluoride membranes and blocked in 1% bovine serum albumin and 0.5% Tween-Tris-buffered saline. Serum samples from 6 horses with **S. neurona** infections (confirmed by culture from neural tissue) and 57 horses without infections (horses from the Eastern Hemisphere, where **S. neurona** does not exist) were tested by Western blot. Horses from both groups had reactivity to the 62-, 30-, 16-, 13-, 11-, 10.5-, and 10-kD bands. Testing was repeated with another step. Blots were **treated** with bovine **S. cruzi** **antibodies** prior to loading the equine samples. After this modification of the Western blot test, positive infection status was significantly associated with reactivity to the 30- and 16-kD bands ( $P < 0.001$ , Fisher's exact test). The **S. cruzi** **antibody**-blocked Western blot had a sample sensitivity of 100% and sample specificity of 98%. It is concluded that the specificity of the Western blot test is improved by blocking proteins not specific to **S. neurona** and using reactivity to the 30- and 16-kD bands as the criterion for a positive test.

L25 ANSWER 16 OF 23 WPIDS (C) 2003 THOMSON DERWENT

AN 1999-571872 [48] WPIDS

DNN N1999-421433 DNC C1999-166894

TI Biologically pure culture of equine Neospora, used as source of vaccines and diagnostic reagents.

DC B04 C06 C07 D16 S03

IN BARR, B C; CONRAD, P A; MARSH, A E

PA (REGC) UNIV CALIFORNIA

CYC 23

PI WO 9947927 A1 19990923 (199948)\* EN 47p

RW: AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL PT SE

W: AU CA JP

AU 9931874 A 19991011 (200008)

US 6071737 A 20000606 (200033)

EP 1064550 A1 20010103 (200102) EN

R: AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE

JP 2002509702 W 20020402 (200225) 47p

ADT WO 9947927 A1 WO 1999-US5754 19990316; AU 9931874 A AU 1999-31874

19990316; US 6071737 A US 1998-42600 19980316; EP 1064550 A1 EP

1999-913906 19990316, WO 1999-US5754 19990316; JP 2002509702 W WO

1999-US5754 19990316, JP 2000-537071 19990316

FDT AU 9931874 A Based on WO 9947927; EP 1064550 A1 Based on WO 9947927; JP

2002509702 W Based on WO 9947927

PRAI US 1998-42600 19980316

AB WO 9947927 A UPAB: 19991122

NOVELTY - Biologically pure culture of equine Neospora, is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the

following:

- (a) detecting **antibodies** (Ab) specifically reactive with equine Neospora antigens (Ag) by forming an Ab-Ag complex;
- (b) detecting Neospora by forming a complex with an **antibody** (Ab1) specifically reactive with Neospora antigen;
- (c) detecting Neospora-specific nucleic acid (I) by hybridization with a specific oligonucleotide probe; and
- (d) pharmaceutical composition containing equine Neospora immunogen and a carrier.

ACTIVITY - Antiprotozoal.

MECHANISM OF ACTION - Induction of a specific immune response.

USE - Immunogens (optionally expressed from gene **therapy** vectors) from equine Neospora are used in vaccines for **treatment** or prevention of Neospora infection in horses and other animals. Neospora is a causative agent of equine protozoal myeloencephalitis (EPM). Detection of Neospora-specific antigens, **antibodies** or nucleic acid (by usual immunoassay or hybridization tests) is used to diagnose infection. **Antibodies** (Ab) specific for equine Neospora are used for diagnosis; to select candidate immunogens for vaccine development; to isolate proteins; to screen DNA libraries and as **therapeutic** /prophylactic agents.

ADVANTAGE - Reagents specific for equine Neospora allow differentiation between equine protozoal myeloencephalitis caused by Neospora and **Sarcocystis neurona**. These pathogens require different **treatments** and **treatment** of Neospora is only effective if applied before the parasite has formed cysts. The vaccines also prevent shedding of oocysts by animals known to be infected.  
Dwg.0/2

L25 ANSWER 17 OF 23 MEDLINE  
AN 2000080043 MEDLINE  
DN 20080043 PubMed ID: 10613219  
TI Encephalomyelitis associated with a **Sarcocystis neurona**-like organism in a sea otter.  
CM Comment in: J Am Vet Med Assoc. 2000 Feb 1;216(3):329  
AU Rosonke B J; Brown S R; Tornquist S J; Snyder S P; Garner M M; Blythe L L  
CS Animal Medical Care of Newport, OR 97365, USA.  
SO JOURNAL OF THE AMERICAN VETERINARY MEDICAL ASSOCIATION, (1999 Dec 15) 215 (12) 1839-42, 1807.  
Journal code: 7503067. ISSN: 0003-1488.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200002  
ED Entered STN: 20000229  
Last Updated on STN: 20000525  
Entered Medline: 20000215  
AB An adult female sea otter housed for 5 years in an outdoor habitat in an aquarium developed signs of neurologic disease. Bilateral caudal paresis was evident initially and other neurologic signs consistent with CNS disease developed rapidly. Diagnostic work-up included CBC, serum biochemical analyses, determination of serum **antibody** titers, radiography of the vertebral column, CSF analysis, muscle biopsy, computed tomography of the brain, and assays for mercury, lead, and thiamine. A tentative diagnosis of encephalitis caused by a **Sarcocystis neurona**-like organism was made on the basis of detection of CSF **antibodies** by use of Western blot analysis. Response to **treatment** was not satisfactory and the sea otter was euthanatized. Immunohistochemical staining revealed S neurona-like organisms within foci of inflammation in the brain and spinal cord. This report provides evidence that, for sea otters, there may be a mode of transmission of an S neurona-like organism that does not involve opossums.



Equine protozoare Myeloenzephalitis bei einem importierten Paint-Horse  
 AU Weigand, K.; Grabner, A.  
 CS Chirurgische Tierklinik der LMU Munchen, Veterinarstr. 13, 80539 Munchen, Germany.  
 SO Pferdeheilkunde, (1997) Vol. 13, No. 3, pp. 231-234. 19 ref.  
 ISSN: 0177-7726  
 DT Journal  
 LA German  
 SL English  
 AB EPM is reported in a 7-year-old American Paint mare, 6 months after it was imported from Florida, USA, to Germany. The mare was referred to the veterinary clinic at Munich University in late pregnancy with severe neurological disorders characterized by ataxia, facial paresis, dullness and considerable problems with uptake of food and water. Mononuclear pleocytosis and elevated protein and lactate concentrations were detected in cerebrospinal fluid (CSF). Diagnosis of active EPM was confirmed by positive Western Blot reactivity on CSF and detection of **antibodies to Sarcocystis neurona**. The condition of the mare improved during 3 weeks of drug **therapy** by trimethoprim sulfonamide and palliative **treatment**. After parturition, the mare and the foal were discharged in good condition. **Treatment** with oral trimethoprim sulfonamide for a minimum of 90 days was continued to avoid relapse of EPM.

L25 ANSWER 22 OF 23 MEDLINE  
 AN 97260250 MEDLINE  
 DN 97260250 PubMed ID: 9106345  
 TI Equine protozoal myeloencephalitis.  
 AU MacKay R J  
 CS Department of Large Animal Clinical Sciences, University of Florida, College of Veterinary Medicine, Gainesville, USA.  
 SO VETERINARY CLINICS OF NORTH AMERICA. EQUINE PRACTICE, (1997 Apr) 13 (1) 79-96. Ref: 72  
 Journal code: 8511904. ISSN: 0749-0739.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 General Review; (REVIEW)  
 (REVIEW, TUTORIAL)  
 LA English  
 FS Priority Journals  
 EM 199706  
 ED Entered STN: 19970612  
 Last Updated on STN: 19970612  
 Entered Medline: 19970603  
 AB Equine protozoal myeloencephalitis (EPM) is a common neurologic disease of horses in the Americas. Horses with EPM most commonly have abnormalities of gait, but they also may present with signs of brain disease. The disease ranges in severity from mild lameness to sudden recumbency, and clinical signs usually are progressive. A causative agent, **Sarcocystis neurona**, has been isolated from affected horses, and serologic surveys suggest that approximately 50% of horses in the United States have been exposed. EPM is considered a **treatable** disease, although the response to antimicrobial **treatment** often is incomplete. This article highlights new information about the life cycle of *S. neurona* and reviews the literature regarding diagnosis, clinical signs, and **treatment** of the disease.

L25 ANSWER 23 OF 23 MEDLINE DUPLICATE 11  
 AN 93385225 MEDLINE  
 DN 93385225 PubMed ID: 8373858  
 TI Immunohistochemical study to demonstrate **Sarcocystis neurona** in equine protozoal myeloencephalitis.

AU Hamir A N; Moser G; Galligan D T; Davis S W; Granstrom D E; Dubey J P  
CS University of Pennsylvania School of Veterinary Medicine, New Bolton  
Center, Kennett Square 19348.  
SO JOURNAL OF VETERINARY DIAGNOSTIC INVESTIGATION, (1993 Jul) 5 (3) 418-22.  
Journal code: 9011490. ISSN: 1040-6387.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199310  
ED Entered STN: 19931105  
Last Updated on STN: 19931105  
Entered Medline: 19931021  
AB A 5-year (1985-1989) retrospective immunohistochemical study was conducted using an avidin-biotin complex (ABC) immunoperoxidase method to demonstrate **Sarcocystis neurona** in histologically suspect cases of equine protozoal myeloencephalitis (EPM). Primary **antibodies** against *S. neurona* and *S. cruzi* were utilized for the ABC technique. The findings were compared with those from cases in which the organisms were detected by examination of hematoxylin and eosin (HE)-stained neuronal sections. HE-stained sections detected the presence of the organisms in 20% of the suspect cases; whereas the ABC technique confirmed the presence of *S. neurona* in 51% and 67% of the cases by *S. neurona* and *S. cruzi* **antibodies**, respectively. A review of clinical case histories showed that 21/47 (45%) of the EPM horses with parasites in the tissue sections had prior **treatment** with antiprotozoal drugs and/or steroids. Using the test results of *S. neurona* and *S. cruzi* as a standard reference, HE test sensitivity based on examination of up to 30 neuronal sections per case was only 25%, and test specificity was 91%.

=> s 19 and sarcocystis neurona  
L10 39 L9 AND SARCOCYSTIS NEURONA

=> s 110 and (16 kd or 30 kd)  
L11 1 L10 AND (16 KD OR 30 KD)

=> d bib ab

L11 ANSWER 1 OF 1 MEDLINE  
AN 2000152631 MEDLINE  
DN 20152631 PubMed ID: 10690772  
TI Improvement of western blot test specificity for detecting equine serum antibodies to **Sarcocystis neurona**.  
AU **Rossano M G; Mansfield L S; Kaneene J B; Murphy A J; Brown C M; Schott H C 2nd; Fox J C**  
CS Animal Health Diagnostic Laboratory, The Population Medicine Center, Michigan State University, East Lansing 48824, USA.  
SO JOURNAL OF VETERINARY DIAGNOSTIC INVESTIGATION, (2000 Jan) 12 (1) 28-32. Journal code: 9011490. ISSN: 1040-6387.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200003  
ED Entered STN: 20000330  
Last Updated on STN: 20000330  
Entered Medline: 20000321  
AB Equine protozoal myeloencephalitis (EPM) is a neurological disease of horses and ponies caused by the apicomplexan protozoan parasite **Sarcocystis neurona**. The purposes of this study were to develop the most stringent criteria possible for a positive test result, to estimate the sensitivity and specificity of the EPM Western blot antibody test, and to assess the ability of bovine antibodies to **Sarcocystis cruzi** to act as a blocking agent to minimize false-positive results in the western blot test for **S. neurona**. **Sarcocystis neurona** merozoites harvested from equine dermal cell culture were heat denatured, and the proteins were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis in a 12-20% linear gradient gel. Separated proteins were electrophoretically transferred to polyvinylidene fluoride membranes and blocked in 1% bovine serum albumin and 0.5% Tween-Tris-buffered saline. Serum samples from 6 horses with **S. neurona** infections (confirmed by culture from neural tissue) and 57 horses without infections (horses from the Eastern Hemisphere, where **S. neurona** does not exist) were tested by Western blot. Horses from both groups had reactivity to the 62-, 30-, 16-, 13-, 11-, 10.5-, and 10-kD bands. Testing was repeated with another step. Blots were treated with bovine **S. cruzi** antibodies prior to loading the equine samples. After this modification of the Western blot test, positive infection status was significantly associated with reactivity to the 30- and 16-kD bands ( $P < 0.001$ , Fisher's exact test). The **S. cruzi** antibody-blocked Western blot had a sample sensitivity of 100% and sample specificity of 98%. It is concluded that the specificity of the Western blot test is improved by blocking proteins not specific to **S. neurona** and using reactivity to the 30- and 16-kD bands as the criterion for a positive test.

=> d his

(FILE 'HOME' ENTERED AT 14:49:07 ON 15 MAY 2003)

FILE 'BIOSIS, MEDLINE, AGRICOLA, EMBASE, CABA, WPIDS, JAPIO, BIOTECHDS, LIFESCI, CAPLUS' ENTERED AT 14:49:30 ON 15 MAY 2003

L1           30 S E2-E3  
               E MANSFIELD L S/AU  
 L2           139 S E3  
               E ROSSANO MARY G/AU  
 L3           9 S E3  
               E ROSSANO M G/AU  
 L4           25 S E3  
               E MURPHY ALICE J/AU  
 L5           15 S E2 OR E3  
               E MURPHY A J/AU  
 L6           337 S E3  
               E VRABLE RUTH A/AU  
 L7           11 S E2 OR E3  
               E VRABLE R A/AU  
 L8           23 S E2 OR E3  
 L9           496 S L1-L8  
 L10          39 S L9 AND SARCOCYSTIS NEURONA  
 L11          1 S L10 AND (16 KD OR 30 KD)

=> s l9 and antibod?

L12           51 L9 AND ANTIBOD?

=> s l10 and antibod?

L13           26 L10 AND ANTIBOD?

=> dup rem l13

PROCESSING COMPLETED FOR L13

L14           12 DUP REM L13 (14 DUPLICATES REMOVED)

=> d bib ab 1-12

L14 ANSWER 1 OF 12 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE  
1

AN 2003:159207 BIOSIS

DN PREV200300159207

TI A herd-level analysis of risk factors for **antibodies** to  
**Sarcocystis neurona** in Michigan equids.

AU **Rossano, M. G.**; Kaneene, J. B. (1); Marteniuk, J. V.; Banks, B.  
D.; Schott, H. C., II; **Mansfield, L. S.**

CS (1) College of Veterinary Medicine, Population Medicine Center, A-109  
Veterinary Medical Center, Michigan State University, East Lansing, MI,  
48824-1314, USA: kaneene@cvm.msu.edu USA

SO Preventive Veterinary Medicine, (15 February 2003) Vol. 57, No. 1-2, pp.  
7-13. print.

ISSN: 0167-5877.

DT Article

LA English

AB Equine protozoal myeloencephalitis (EPM) is a neurological disease of horses and ponies caused by infection of the central nervous system with the protozoan parasite **Sarcocystis neurona**. A herd-level analysis of a cross-sectional study of serum **antibodies** to *S. neurona* in Michigan equids was conducted, using data collected in 1997 for study that included 1121 equids from 98 Michigan horse farms. Our objective was to identify specific herd-level risk factors associated with seropositivity. We tested associations between herd seroprevalence and various farm-management practices (including feed-storage methods and wildlife control). Multivariable models were developed for three strata based on relative opossum abundance (opossum districts). Herd seroprevalence ranged from 0 to 100% (median = 57%); No risk factor was significantly associated with herd seroprevalence at  $P \leq 0.05$  in all opossum districts. Our results suggest that equids living in areas with large opossum populations might be infected with *S. neurona* from multiple

sources.

L14 ANSWER 2 OF 12 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
AN 2003:67219 BIOSIS  
DN PREV200300067219  
TI Immunoassay for equine protozoal myeloencephalitis in horses.  
AU **Mansfield, Linda S.; Murphy, Alice J. (1);**  
**Rossano, Mary G.**  
CS (1) St. Johns, MI, USA USA  
ASSIGNEE: Board of Trustees of Michigan State University  
PI US 6489148 December 03, 2002  
SO Official Gazette of the United States Patent and Trademark Office Patents,  
(Dec. 3 2002) Vol. 1265, No. 1, pp. No Pagination.  
<http://www.uspto.gov/web/menu/patdata.html>. e-file.  
ISSN: 0098-1133.  
DT Patent  
LA English  
AB An immunoassay for *Sarcocystis* neurons **antibodies** in equines is described. The immunoassay uses blocking of *Sarcocystis* antigens by **antibodies** to *Sarcocystis* sp. other than **Sarcocystis neurona** in connection with the immunoassay.

L14 ANSWER 3 OF 12 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
AN 2002:234586 BIOSIS  
DN PREV200200234586  
TI Antigen test to detect equine protozoal myeloencephalitis in horse serum and cerebrospinal fluid.  
AU **Mansfield, Linda S.; Rossano, Mary G.; Murphy,**  
**Alice J.; Vrable, Ruth A. (1)**  
CS (1) Williamston, MI USA  
ASSIGNEE: Board of Trustees of Michigan State University, East Lansing, MI, USA  
PI US 6344337 February 05, 2002  
SO Official Gazette of the United States Patent and Trademark Office Patents,  
(Feb. 5, 2002) Vol. 1255, No. 1, pp. No Pagination.  
<http://www.uspto.gov/web/menu/patdata.html>. e-file.  
ISSN: 0098-1133.  
DT Patent  
LA English  
AB The present invention provides an immunoassay to detect identifying antigens in horses that are infected with **Sarcocystis neurona**. The immunoassay is preferably an antigen-capture-based assay that relies upon polyclonal or monoclonal **antibodies** against a 16 (+-4) and/or 30 (+-4) kDa antigens specific to **Sarcocystis neurona** to detect the presence of the 16 (+-4) and/or 30 (+-4) kDa antigens in equine serum or equine cerebrospinal fluid.

L14 ANSWER 4 OF 12 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE  
2  
AN 2002:556058 BIOSIS  
DN PREV200200556058  
TI Cross-sectional study of serum **antibodies** against **Sarcocystis neurona** in cats tested for **antibodies** against *Toxoplasma gondii*.  
AU **Rossano, Mary G.; Murphy, Alice J.; Vrable, Ruth**  
**A.; Vanzo, Nicole E.; Lewis, Stacy K.; Sheline, Katherine D.;**  
**Kaneene, John B.; Mansfield, Linda S. (1)**  
CS (1) Animal Health Diagnostic Laboratory, College of Veterinary Medicine,  
Michigan State University, East Lansing, MI, 48824 USA  
SO Journal of the American Veterinary Medical Association, (August 15, 2002)  
Vol. 221, No. 4, pp. 511-514. <http://www.avma.org>. print.  
ISSN: 0003-1488.

MECHANISM OF ACTION - Vaccine.

USE - The vaccines and methods are used for protecting equids against infections by the protozoan parasite *Sarcocystis neurona*.  
Dwg.0/0

- L14 ANSWER 6 OF 12 BIOTECHDS COPYRIGHT 2003 THOMSON DERWENT AND ISI  
AN 2001-07735 BIOTECHDS  
TI Vaccinating equids against protozoal **Sarcocystis neurona** infections using antigens;  
**Sarcocystis neurona** nucleic acid vaccine and recombinant vaccine  
AU **Mansfield L S; Rossano M G; Murphy A J; Vrable R A**  
PA Univ.Michigan-State  
LO East Lansing, MI, USA.  
PI WO 2001015708 8 Mar 2001  
AI WO 2000-US24221 31 Aug 2000  
PRAI US 2000-513086 24 Feb 2000; US 1999-152193 2 Sep 1999  
DT Patent  
LA English  
OS WPI: 2001-218486 [22]  
AB A method for vaccinating equids against **Sarcocystis neurona** infection is claimed. It involves using protein groups of unique 16(+4) or 30(+4) antigens of *S. neurona*. Also claimed are: a vaccine (I) for providing passive immunity to **Sarcocystis neurona** infection; a vaccine (II) for active immunization of an equid against a *S. neurona* infection; a vaccine (III) for protecting an equid from *S. neurona* infection; a method (IV or V) for vaccinating an equid against a *S. neurona* infection; a method (VI) of providing passive immunity to a *S. neurona* infection; a method (VII) for producing a protein (e.g. glutathione-transferase); a method (VIII) for producing an **antibody**; providing a microorganism in a culture containing DNA encoding a fusion protein; a monoclonal **antibody** (IX); an isolated DNA (X); a bacterial clone (XI); a vaccine (XII) for an equid containing an isolated recombinant protein; a vaccine (XIII or XIV) for an equid containing recombinant virus vector containing DNA; and a method (XV) for protecting an equid against *S. neurona*. The vaccines and methods are used for protecting equids against infection by the protozoan parasite *Sarcocystis neurona*. (54pp)
- L14 ANSWER 7 OF 12 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE  
4  
AN 2001:135932 BIOSIS  
DN PREV200100135932  
TI The seroprevalence of **antibodies** to **Sarcocystis neurona** in Michigan equids.  
AU **Rossano, M. G.**; Kaneene, J. B. (1); Marteniuk, J. V.; Banks, B. D.; Schott, H. C., II; **Mansfield, L. S.**  
CS (1) Population Medicine Center, College of Veterinary Medicine, A-109 Veterinary Medical Center, Michigan State University, East Lansing, MI, 48824-1314: kaneene@cvm.msu.edu USA  
SO Preventive Veterinary Medicine, (29 January, 2001) Vol. 48, No. 2, pp. 113-128. print.  
ISSN: 0167-5877.  
DT Article  
LA English  
SL English  
AB A cross-sectional study of serum **antibodies** to **Sarcocystis neurona** (the etiologic agent of equine protozoal myeloencephalitis, EPM) was performed on Michigan equids. Our objectives were to determine the seroprevalence of **antibodies** to *S. neurona* in Michigan equids and to identify specific risk factors for seropositivity. A random, weighted sample of Michigan horse farms

conjugate and then detected by reaction with an appropriate color forming substrate. The **antibody** is immobilized on a support chosen from a membrane or a plate. (64pp)

L14 ANSWER 12 OF 12 MEDLINE DUPLICATE 7  
AN 2000152631 MEDLINE  
DN 20152631 PubMed ID: 10690772  
TI Improvement of western blot test specificity for detecting equine serum **antibodies** to **Sarcocystis neurona**.  
AU **Rossano M G; Mansfield L S; Kaneene J B; Murphy A**  
J; Brown C M; Schott H C 2nd; Fox J C  
CS Animal Health Diagnostic Laboratory, The Population Medicine Center,  
Michigan State University, East Lansing 48824, USA.  
SO JOURNAL OF VETERINARY DIAGNOSTIC INVESTIGATION, (2000 Jan) 12 (1) 28-32.  
Journal code: 9011490. ISSN: 1040-6387.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200003  
ED Entered STN: 20000330  
Last Updated on STN: 20000330  
Entered Medline: 20000321  
AB Equine protozoal myeloencephalitis (EPM) is a neurological disease of horses and ponies caused by the apicomplexan protozoan parasite **Sarcocystis neurona**. The purposes of this study were to develop the most stringent criteria possible for a positive test result, to estimate the sensitivity and specificity of the EPM Western blot **antibody** test, and to assess the ability of bovine **antibodies** to **Sarcocystis cruzi** to act as a blocking agent to minimize false-positive results in the western blot test for **S. neurona**. **Sarcocystis neurona** merozoites harvested from equine dermal cell culture were heat denatured, and the proteins were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis in a 12-20% linear gradient gel. Separated proteins were electrophoretically transferred to polyvinylidene fluoride membranes and blocked in 1% bovine serum albumin and 0.5% Tween-Tris-buffered saline. Serum samples from 6 horses with **S. neurona** infections (confirmed by culture from neural tissue) and 57 horses without infections (horses from the Eastern Hemisphere, where **S. neurona** does not exist) were tested by Western blot. Horses from both groups had reactivity to the 62-, 30-, 16-, 13-, 11-, 10.5-, and 10-kD bands. Testing was repeated with another step. Blots were treated with bovine **S. cruzi** **antibodies** prior to loading the equine samples. After this modification of the Western blot test, positive infection status was significantly associated with reactivity to the 30- and 16-kD bands ( $P < 0.001$ , Fisher's exact test). The **S. cruzi** **antibody**-blocked Western blot had a sample sensitivity of 100% and sample specificity of 98%. It is concluded that the specificity of the Western blot test is improved by blocking proteins not specific to **S. neurona** and using reactivity to the 30- and 16-kD bands as the criterion for a positive test.

=> d his

(FILE 'HOME' ENTERED AT 14:49:07 ON 15 MAY 2003)

FILE 'BIOSIS, MEDLINE, AGRICOLA, EMBASE, CABA, WPIDS, JAPIO, BIOTECHDS, LIFESCI, CAPLUS' ENTERED AT 14:49:30 ON 15 MAY 2003

E MANSFIELD LINDA S/AU  
L1 30 S E2-E3  
E MANSFIELD L S/AU  
L2 139 S E3

Rossano, Mary G.

CS (1) St. Johns, MI, USA USA  
ASSIGNEE: Board of Trustees of Michigan State University  
PI US 6489148 December 03, 2002  
SO Official Gazette of the United States Patent and Trademark Office Patents,  
(Dec. 3 2002) Vol. 1265, No. 1, pp. No Pagination.  
<http://www.uspto.gov/web/menu/patdata.html>. e-file.  
ISSN: 0098-1133.  
DT Patent  
LA English  
AB An immunoassay for *Sarcocystis* neurons antibodies in equines is described.  
The immunoassay uses blocking of *Sarcocystis* antigens by antibodies to  
*Sarcocystis* sp. other than *Sarcocystis neurona* in  
connection with the immunoassay.

L15 ANSWER 3 OF 15 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
AN 2002:234586 BIOSIS  
DN PREV200200234586  
TI Antigen test to detect equine protozoal myeloencephalitis in horse serum  
and cerebrospinal fluid.  
AU Mansfield, Linda S.; Rossano, Mary G.; Murphy,  
Alice J.; Vrable, Ruth A. (1)  
CS (1) Williamston, MI USA  
ASSIGNEE: Board of Trustees of Michigan State University, East Lansing,  
MI, USA  
PI US 6344337 February 05, 2002  
SO Official Gazette of the United States Patent and Trademark Office Patents,  
(Feb. 5, 2002) Vol. 1255, No. 1, pp. No Pagination.  
<http://www.uspto.gov/web/menu/patdata.html>. e-file.  
ISSN: 0098-1133.  
DT Patent  
LA English  
AB The present invention provides an immunoassay to detect identifying  
antigens in horses that are infected with *Sarcocystis*  
*neurona*. The immunoassay is preferably an antigen-capture-based  
assay that relies upon polyclonal or monoclonal antibodies against a 16  
(+/-4) and/or 30 (+/-4) kDa antigens specific to *Sarcocystis*  
*neurona* to detect the presence of the 16 (+/-4) and/or 30 (+/-4) kDa  
antigens in equine serum or equine cerebrospinal fluid.

L15 ANSWER 4 OF 15 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE  
2  
AN 2002:556058 BIOSIS  
DN PREV200200556058  
TI Cross-sectional study of serum antibodies against *Sarcocystis*  
*neurona* in cats tested for antibodies against *Toxoplasma gondii*.  
AU Rossano, Mary G.; Murphy, Alice J.; Vrable, Ruth  
A.; Vanzo, Nicole E.; Lewis, Stacy K.; Sheline, Katherine D.;  
Kaneene, John B.; Mansfield, Linda S. (1)  
CS (1) Animal Health Diagnostic Laboratory, College of Veterinary Medicine,  
Michigan State University, East Lansing, MI, 48824 USA  
SO Journal of the American Veterinary Medical Association, (August 15, 2002)  
Vol. 221, No. 4, pp. 511-514. <http://www.avma.org>. print.  
ISSN: 0003-1488.  
DT Article  
LA English  
AB Objective-To determine apparent seroprevalence of antibodies against  
*Sarcocystis neurona* in a population of domestic cats  
previously tested for antibodies against *Toxoplasma gondii*.  
Design-Cross-sectional study. Sample Population-Serum from 196 domestic  
cats. Procedure-Banked serum samples submitted to the Michigan State  
University Animal Health Diagnostic Laboratory for *T gondii* diagnostic  
testing were tested for antibodies against *S neurona* by use of an indirect



reacted with peroxidase conjugate and then detected by reaction with an appropriate color forming substrate. The antibody is immobilized on a support chosen from a membrane or a plate. (64pp)

L15 ANSWER 14 OF 15 MEDLINE DUPLICATE 8  
AN 2000152631 MEDLINE  
DN 20152631 PubMed ID: 10690772  
TI Improvement of western blot test specificity for detecting equine serum antibodies to **Sarcocystis neurona**.  
AU **Rossano M G; Mansfield L S; Kaneene J B; Murphy A**  
J; Brown C M; Schott H C 2nd; Fox J C  
CS Animal Health Diagnostic Laboratory, The Population Medicine Center, Michigan State University, East Lansing 48824, USA.  
SO JOURNAL OF VETERINARY DIAGNOSTIC INVESTIGATION, (2000 Jan) 12 (1) 28-32.  
Journal code: 9011490. ISSN: 1040-6387.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200003  
ED Entered STN: 20000330  
Last Updated on STN: 20000330  
Entered Medline: 20000321  
AB Equine protozoal myeloencephalitis (EPM) is a neurological disease of horses and ponies caused by the apicomplexan protozoan parasite **Sarcocystis neurona**. The purposes of this study were to develop the most stringent criteria possible for a positive test result, to estimate the sensitivity and specificity of the EPM Western blot antibody test, and to assess the ability of bovine antibodies to **Sarcocystis cruzi** to act as a blocking agent to minimize false-positive results in the western blot test for **S. neurona**. **Sarcocystis neurona** merozoites harvested from equine dermal cell culture were heat denatured, and the proteins were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis in a 12-20% linear gradient gel. Separated proteins were electrophoretically transferred to polyvinylidene fluoride membranes and blocked in 1% bovine serum albumin and 0.5% Tween-Tris-buffered saline. Serum samples from 6 horses with **S. neurona** infections (confirmed by culture from neural tissue) and 57 horses without infections (horses from the Eastern Hemisphere, where **S. neurona** does not exist) were tested by Western blot. Horses from both groups had reactivity to the 62-, 30-, 16-, 13-, 11-, 10.5-, and 10-kD bands. Testing was repeated with another step. Blots were treated with bovine **S. cruzi** antibodies prior to loading the equine samples. After this modification of the Western blot test, positive infection status was significantly associated with reactivity to the 30- and 16-kD bands ( $P < 0.001$ , Fisher's exact test). The **S. cruzi** antibody-blocked Western blot had a sample sensitivity of 100% and sample specificity of 98%. It is concluded that the specificity of the Western blot test is improved by blocking proteins not specific to **S. neurona** and using reactivity to the 30- and 16-kD bands as the criterion for a positive test.

L15 ANSWER 15 OF 15 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC. DUPLICATE 9  
AN 2000:21537 BIOSIS  
DN PREV200000021537  
TI Simplified technique for isolation, excystation, and culture of **Sarcocystis** species from opossums.  
AU **Murphy, A. J. (1); Mansfield, L. S.**  
CS (1) Animal Health Diagnostic Laboratory, Michigan State University, East Lansing, MI, 48824 USA  
SO Journal of Parasitology, (Oct., 1999) Vol. 85, No. 5, pp. 979-981.  
ISSN: 0022-3395.  
DT Article

and analyzed both qualitatively (western blot) and quantitatively (anti-17kDa) for anti-S. neurona IgG. Four of the challenged horses were given dexamethasone (0.1mg/kg orally once daily) for the duration of the experiment. All challenged horses immunoconverted against S. neurona in blood within 32 days of challenge and in CSF within 61 days. There was a trend ( $P = 0.057$ ) for horses given dexamethasone to immunoconvert earlier than horses that were not immunosuppressed. Anti-17kDa was detected in the CSF of all challenged horses by day 61. This response was statistically greater at day 32 in horses given dexamethasone. Control horses remained seronegative throughout the period in which all challenged horses converted. One control horse immunoconverted in blood at day 75 and in CSF at day 89. Signs of neurologic disease were mild to equivocal in challenged horses. Horses given dexamethasone had more severe signs of limb weakness than did horses not given dexamethasone; however, we could not determine whether these signs were due to spinal cord disease or to effects of systemic illness. At necropsy, mild-moderate multifocal gliosis and neurophagia were found histologically in the spinal cords of 7/8 challenged horses. No organisms were seen either in routinely processed sections or by immunohistochemistry. Although neurologic disease comparable to naturally occurring equine protozoal myeloencephalitis (EPM) was not produced, we had clear evidence of an immune response to challenge both systemically and in the CNS. Broad immunosuppression with dexamethasone did not increase the severity of histologic changes in the CNS of challenged horses. Future work must focus on defining the factors that govern progression of inapparent S. neurona infection to EPM.

L20 ANSWER 3 OF 3 MEDLINE  
 AN 93222344 MEDLINE  
 DN 93222344 PubMed ID: 8466988  
 TI Equine protozoal myeloencephalitis: antigen analysis of cultured **Sarcocystis neurona** merozoites.  
 AU Granstrom D E; Dubey J P; Davis S W; Fayer R; Fox J C; Poonacha K B; Giles R C; Comer P F  
 CS Department of Veterinary Science, University of Kentucky, Lexington 40546-0099.  
 SO JOURNAL OF VETERINARY DIAGNOSTIC INVESTIGATION, (1993 Jan) 5 (1) 88-90. Journal code: 9011490. ISSN: 1040-6387.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199305  
 ED Entered STN: 19930521  
 Last Updated on STN: 19930521  
 Entered Medline: 19930510  
 AB Antigens of cultured **Sarcocystis neurona** merozoites were examined using immunoblot analysis. Blotted proteins were probed with S. cruzi, S. muris, and S. neurona antisera produced in rabbits, S. fayeri (pre- and post-infection) and S. neurona (pre- and post-inoculation) sera produced in horses, immune sera from 7 histologically confirmed cases of equine protozoal myeloencephalitis (EPM), and pre-suckle serum from a newborn foal. Eight proteins, 70, 24, 23.5, 22.5, 13, 11, 10.5, and 10 Kd, were detected only by S. neurona antiserum and/or immune serum from EPM-affected horses. Equine sera were titrated by the indirect immunofluorescent **antibody** (IFA) method using air-dried, cultured S. neurona merozoites. Anti-Sarcocystis IFA titers were found in horses with or without EPM. Serum titers did not correspond to the number of specific bands recognized on immunoblots.

=> s 119 and kd

L21 1 L19 AND KD

Entered Medline: 20020117

AB A two-month-old Appaloosa colt developed neurological signs shortly after birth involving deficits affecting cranial nerves IV, VII, VIII, IX, X and XII, and possibly nerve VI. The most likely differential diagnoses were congenital anomalies, meningoencephalitis, trauma or nutritional causes. The foal was investigated by the analysis of cerebrospinal fluid (CSF), electromyography (EMG), brain auditory evoked responses, magnetic resonance imaging (MRI), peripheral nerve biopsy, and Western blot analysis for the presence of intrathecal **antibodies** to **Sarcocystis neurona**, the causative agent of equine protozoal myeloencephalitis. Significantly abnormal EMG findings included spontaneous electrical activity of the tongue, suggesting denervation. The MRI was useful in ruling out masses, congenital anomalies and focal abscessation. The cytology of CSF revealed mild mononuclear reactivity. Western blot testing of CSF was positive, indicating the intrathecal presence of **antibodies** to **S. neurona**. The foal was **treated** with pyrimethamine and trimethoprim-sulphadiazine for two months and returned to nearly normal neurologic status.

L25 ANSWER 10 OF 23 CABA COPYRIGHT 2003 CABI

AN 2001:140175 CABA

DN 20013139273

TI Efficacy of ponazuril 15% oral paste as a **treatment** for equine protozoal myeloencephalitis

AU Furr, M.; Kennedy, T.; MacKay, R.; Reed, S.; Andrews, F.; Bernard, B.; Bain, F.; Byars, D.

CS Virginia-Maryland Regional College of Veterinary Medicine, Marion duPont Scott Equine Medical Center, PO Box 1938, Leesburg, VA 20177, USA.

SO Veterinary Therapeutics, (2001) Vol. 2, No. 3, pp. 215-222. 15 ref. ISSN: 1528-3593

DT Journal

LA English

AB Equine protozoal myeloencephalitis (EPM) is a neurologic disease of horses most commonly caused by the protozoan parasite **Sarcocystis neurona**. Until recently the only **treatment** option was the combination of a sulfonamide with pyrimethamine. The present study was performed to assess the efficacy of ponazuril, an anticoccidial triazine-based compound, as a **treatment** for naturally occurring EPM. 101 horses with EPM were randomly allocated to **treatment** with ponazuril 15% oral paste at either 5 or 10 mg/kg body weight for 28 consecutive days. Horses were evaluated clinically and by analysis of blood and cerebrospinal fluid (CSF) before and 28 and 118 days after the start of **treatment**. Clinical success was defined as either an improvement in neurologic score by at least one grade (on a 0 to 5 scale) or conversion to negative status on Western blot for **S. neurona antibodies** by 20 days following cessation of **treatment**. Overall, 62% of the horses, including 28 of 47 **treated** with ponazuril at 5 mg/kg and 35 of 54 **treated** with 10 mg/kg, met the criteria for successful **treatment**. The Western blot for CSF became negative in 10% (10/101) of the horses. Quantification of the anti-17kDa **antibody** response in Western blot (relative quantity CSF) did not reveal a significant change in response to **treatment**. However, immunoglobulin index did decrease significantly during **treatment** (P=.01). The findings of this study support the efficacy of ponazuril for the **treatment** of EPM.

L25 ANSWER 11 OF 23 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE 4

AN 2001:169888 BIOSIS

DN PREV200100169888

TI Utilization of stress in the development of an equine model for equine protozoal myeloencephalitis.

AU Saville, W. J. A. (1); Stich, R. W.; Reed, S. M.; Njoku, C. J.; Oglesbee,

M. J.; Wunschmann, A.; Grover, D. L.; Larew-Nauble, A. L.; Stanek, J. F.; Granstrom, D. E.; Dubey, J. P.

CS (1) Department of Veterinary Preventive Medicine, College of Veterinary Medicine, Ohio State University, Columbus, OH, 43210: saville.4@osu.edu USA

SO Veterinary Parasitology, (26 February, 2001) Vol. 95, No. 2-4, pp. 211-222. print.  
ISSN: 0304-4017.

DT Article

LA English

SL English

AB Neurologic disease in horses caused by **Sarcocystis**

**neurona** is difficult to diagnose, **treat**, or prevent, due to the lack of knowledge about the pathogenesis of the disease. This in turn is confounded by the lack of a reliable equine model of equine protozoal myeloencephalitis (EPM). Epidemiologic studies have implicated stress as a risk factor for this disease, thus, the role of transport stress was evaluated for incorporation into an equine model for EPM. Sporocysts from feral opossums were bioassayed in interferon-gamma gene knockout (KO) mice to determine minimum number of viable *S. neurona* sporocysts in the inoculum. A minimum of 80,000 viable *S. neurona* sporocysts were fed to each of the nine horses. A total of 12 *S. neurona* **antibody** negative horses were divided into four groups (1-4). Three horses (group 1) were fed sporocysts on the day of arrival at the study site, three horses were fed sporocysts 14 days after acclimatization (group 2), three horses were given sporocysts and dexamethasone 14 days after acclimatization (group 3) and three horses were controls (group 4). All horses fed sporocysts in the study developed **antibodies** to *S. neurona* in serum and cerebrospinal fluid (CSF) and developed clinical signs of neurologic disease. The most severe clinical signs were in horses in group 1 subjected to transport stress. The least severe neurologic signs were in horses **treated** with dexamethasone (group 3). Clinical signs improved in four horses from two **treatment** groups by the time of euthanasia (group 1, day 44; group 3, day 47). Post-mortem examinations, and tissues that were collected for light microscopy, immunohistochemistry, tissue cultures, and bioassay in KO mice, revealed no direct evidence of *S. neurona* infection. However, there were lesions compatible with *S. neurona* infection in horses. The results of this investigation suggest that stress can play a role in the pathogenesis of EPM. There is also evidence to suggest that horses in nature may clear the organism routinely, which may explain the relatively high number of normal horses with CSF **antibodies** to *S. neurona* compared to the prevalence of EPM.

L25 ANSWER 12 OF 23 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE  
5

AN 2001:169887 BIOSIS

DN PREV200100169887

TI Immunoconversion against **Sarcocystis neurona** in normal and dexamethasone-**treated** horses challenged with *S. neurona* sporocysts.

AU Cutler, Tim J.; MacKay, Robert J. (1); Ginn, Pamela E.; Gillis, Karen; Tanhauser, Susan M.; LeRay, Erin V.; Dame, John B.; Greiner, Ellis C.

CS (1) Large Animal Clinical Sciences, College of Veterinary Medicine, University of Florida, Gainesville, FL, 32610: mackayr@mail.vetmed.ufl.edu USA

SO Veterinary Parasitology, (26 February, 2001) Vol. 95, No. 2-4, pp. 197-210. print.  
ISSN: 0304-4017.

DT Article

LA English

SL English

AB Equine protozoal myeloencephalitis is a common neurologic disease of

horses in the Americas usually caused by **Sarcocystis neurona**. To date, the disease has not been induced in horses using characterized sporocysts from *Didelphis virginiana*, the definitive host. *S. neurona* sporocysts from 15 naturally infected opossums were fed to horses seronegative for **antibodies** against *S. neurona*. Eight horses were given  $5 \times 10^5$  sporocysts daily for 7 days. Horses were examined for abnormal clinical signs, and blood and cerebrospinal fluid were harvested at intervals for 90 days after the first day of challenge and analyzed both qualitatively (western blot) and quantitatively (anti-17 kDa) for anti-*S. neurona* IgG. Four of the challenged horses were given dexamethasone (0.1 mg/kg orally once daily) for the duration of the experiment. All challenged horses immunoconverted against *S. neurona* in blood within 32 days of challenge and in CSF within 61 days. There was a trend ( $P = 0.057$ ) for horses given dexamethasone to immunoconvert earlier than horses that were not immunosuppressed. Anti-17 kDa was detected in the CSF of all challenged horses by day 61. This response was statistically greater at day 32 in horses given dexamethasone. Control horses remained seronegative throughout the period in which all challenged horses converted. One control horse immunoconverted in blood at day 75 and in CSF at day 89. Signs of neurologic disease were mild to equivocal in challenged horses. Horses given dexamethasone had more severe signs of limb weakness than did horses not given dexamethasone; however, we could not determine whether these signs were due to spinal cord disease or to effects of systemic illness. At necropsy, mild-moderate multifocal gliosis and neurophagia were found histologically in the spinal cords of 7/8 challenged horses. No organisms were seen either in routinely processed sections or by immunohistochemistry. Although neurologic disease comparable to naturally occurring equine protozoal myeloencephalitis (EPM) was not produced, we had clear evidence of an immune response to challenge both systemically and in the CNS. Broad immunosuppression with dexamethasone did not increase the severity of histologic changes in the CNS of challenged horses. Future work must focus on defining the factors that govern progression of inapparent *S. neurona* infection to EPM.

L25 ANSWER 13 OF 23 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE  
6  
AN 2000:509421 BIOSIS  
DN PREV200000509421  
TI Evaluation of risk factors associated with clinical improvement and survival of horses with equine protozoal myeloencephalitis.  
AU Saville, William J. (1); Morley, Paul S. (1); Reed, Stephen M.; Granstrom, David E.; Kohn, Catherine W.; Hinchcliff, Kenneth W.; Wittum, Thomas E. (1)  
CS (1) Department of Veterinary Preventive Medicine, College of Veterinary Medicine, Ohio State University, Columbus, OH, 43210 USA  
SO Journal of the American Veterinary Medical Association, (October 15, 2000) Vol. 217, No. 8, pp. 1181-1185. print.  
ISSN: 0003-1488.  
DT Article  
LA English  
SL English  
AB Objective: To investigate risk factors for use in predicting clinical improvement and survival of horses with equine protozoal myeloencephalitis (EPM). Design: Longitudinal epidemiologic study. Animals: 251 horses with EPM. Procedure: Between 1992 and 1995, 251 horses with EPM were admitted to our facility. A diagnosis of EPM was made on the basis of neurologic abnormalities and detection of **antibody** to **Sarcocystis neurona** or *S. neurona* DNA in CSF. Data were obtained from hospital records and through telephone follow-up interviews. Factors associated with clinical improvement and survival were analyzed, using multivariable logistic regression. Results: The likelihood of clinical improvement after diagnosis of EPM was lower in horses used for breeding and pleasure activities. **Treatment** for EPM increased the probability that a

L25 ANSWER 18 OF 23 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE  
 9  
 AN 1999:283751 BIOSIS  
 DN PREV199900283751  
 TI Initial experiences with the use of nitazoxanide in the **treatment**  
 of equine protozoal encephalitis in Northern California.  
 AU Vatistas, Nicholas (1); Fenger, Clara; Palma, Kathleen; Sifferman, Roger  
 CS (1) VM Surgical and Radiological Sciences, University of California,  
 Tupper Hall, Room 2112, Davis, CA, 95688 USA  
 SO Equine Practice, (May, 1999) Vol. 21, No. 5, pp. 18-21.  
 ISSN: 0162-8941.  
 DT Article  
 LA English  
 SL English  
 AB Equine protozoal myeloencephalitis (EPM) is the most common neurological  
 condition affecting horses in North and South America. Nitazoxanide has  
 been reported to be effective against a wide variety of parasites and  
 bacteria in both animals and humans, including protozoa, nematodes,  
 cestodes, trematodes, almost all anaerobic obligate and facultative  
 bacteria, and some aerobic bacteria. The purpose of this study was to  
 determine the effectiveness of nitazoxanide for the **treatment** of  
 EPM in horses. For inclusion in the study, horses had to have evidence of  
 proprioceptive deficits in one or more limbs, and have a positive  
 immunoblot (Western blot) assay for **Sarcocystis neurona**  
**antibodies** in cerebrospinal fluid. The degree of ataxia was graded  
 from 0 (none) to 5 (severe). Seven horses fit the criteria for inclusion,  
 five horses were grade 2, one horse was a grade 3, and one horse was a  
 grade 4. Nitazoxanide was administered as a feed additive, as tablets, as  
 a powder, or as a paste at 50 or 75 mg/kg for approximately 28 days. Two  
 horses became inappetent and depressed during the course of  
**treatment**. However, no long-term sequelae were noted. Four horses  
 became pregnant while on the medication, and remained pregnant at the end  
 of the study period. Neurologic signs returned in two horses, and  
 medication was re-introduced. By the end of the trial (85 to 140 days),  
 five horses were neurologically normal, one horse had improved from a  
 grade 4 to a grade 1, and one horse was unchanged. Cerebrospinal fluid  
 samples were obtained from approximately 85 to 140 days after the start of  
 medication. The samples remained positive for **Sarcocystis**  
**neurona antibodies** by immunoblot (Western blot).  
 However, in six of the seven horses, the relative quantity of  
**antibody** had decreased. In its final formulation as a paste,  
 nitazoxanide was well accepted and well tolerated by horses. It improved  
 the neurological status of six of the seven horses. Nitazoxanide has the  
 advantage over presently available medications in that it is cidal (in  
 other species) rather than static in action, it has been administered to  
 pregnant rodents without inducing signs of fetal abnormalities, and is  
 available in a formulation that is more easily administered to horses.

L25 ANSWER 19 OF 23 MEDLINE  
 AN 1999048790 MEDLINE  
 DN 99048790 PubMed ID: 9831950  
 TI Neospora caninum-associated equine protozoal myeloencephalitis.  
 AU Hamir A N; Tornquist S J; Gerros T C; Topper M J; Dubey J P  
 CS College of Veterinary Medicine, Oregon State University, Corvallis 97331,  
 USA.  
 SO VETERINARY PARASITOLOGY, (1998 Nov 27) 79 (4) 269-74.  
 Journal code: 7602745. ISSN: 0304-4017.  
 CY Netherlands  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199901

ED Entered STN: 19990202  
Last Updated on STN: 19990202  
Entered Medline: 19990119  
AB Equine protozoal myeloencephalitis (EPM) was clinically diagnosed in a 20-year-old horse with severe ataxia. The cerebrospinal fluid was positive for **Sarcocystis neurona antibodies** by western blot. The horse was administered corticosteroids to facilitate in vitro culture of *S. neurona* from its spinal cord following necropsy. Microscopic lesions of EPM were present in the brain and in the spinal cord, including multifocal inflammatory cellular infiltrates and several large groups of protozoa. Immunohistochemical, and light and electron microscopic examinations revealed that the protozoa were *Neospora caninum* and not *S. neurona*. The protozoa divided by endodyogeny, tachyzoites had rhoptries, and organisms reacted specifically to *N. caninum antibodies*. Veterinarians should be aware of increasing diagnosis of *N. caninum* as another etiological agent responsible for the lesions of EPM.

L25 ANSWER 20 OF 23 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE  
10

AN 1997:215661 BIOSIS

DN PREV199799522165

TI Epizootic of equine protozoal myeloencephalitis on a farm.

AU Fenger, Clara K. (1); Granstrom, David E.; Langemeier, John L.; Stamper, Shelby

CS (1) Equine Intern. Med. Consulting, 3288 Valhalla Dr., Lexington, KY 40515  
USA

SO Journal of the American Veterinary Medical Association, (1997) Vol. 210,  
No. 7, pp. 923-927.  
ISSN: 0003-1488.

DT Article

LA English

AB Objective-To determine the clinical findings, course of **treatment**, and long-term outcome of horses on a farm in central Kentucky during an epizootic of equine protozoal myeloencephalitis (EPM). Design-Cohort study. Animals-21 horses on a farm in central Kentucky, 12 of which developed clinical signs of EPM. Procedure-Horses on the farm were serially examined for signs of neurologic disease and serum and CSF **antibodies to Sarcocystis neurona**. Horses were considered to have EPM if they had neurologic signs and positive test results for **antibodies to S neurona** in CSF. Blood values were monitored for evidence of abnormalities resulting from long-term pyrimethamine and trimethoprim-sulfamethoxazole administration. Physical, neurologic, and fetal necropsy examinations were performed as needed. Horses were **treated** for EPM until they had negative test results for CSF **antibodies to S neurona**. Results-Of 21 horses on the farm, 12 had EPM over the course of 6 months. The duration of **treatment** ranged from 45 to 211 days, excluding 1 horse that persistently had CSF **antibodies to S neurona**. Adverse effects from pyrimethamine and trimethoprim-sulfamethoxazole administration included transient fever, anorexia, and depression (n = 2); acute worsening of ataxia (2); mild anemia (4); and abortions (3). Clinical Implications-EPM may develop as an epizootic. In the horses of this report, subtle clinical signs that were originally considered unimportant ultimately progressed to obvious neurologic signs. Adverse effects associated with EPM **treatment** included worsening of neurologic signs, anemia, abortion, and leukopenic and febrile episodes.

L25 ANSWER 21 OF 23 CABA COPYRIGHT 2003 CABI

AN 1998:76244 CABA

DN 982206910

TI Equine protozoal myeloencephalitis (EPM) in an imported American Paint horse

## ANSWER 2 OF 2 MEDLINE

AN 96342244 MEDLINE  
DN 96342244 PubMed ID: 8720045  
TI Toxicity and properties of the extract from **Sarcocystis** cruzi  
cysts.  
AU Saito M; Taguchi K; Shibata Y; Kobayashi T; Shimura K; Itagaki H  
CS Kumagaya Meat Inspection Center Saitama Prefecture, Japan.  
SO JOURNAL OF VETERINARY MEDICAL SCIENCE, (1995 Dec) 57 (6) 1049-51.  
Journal code: A27; 9105360. ISSN: 0916-7250.  
CY Japan  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199610  
ED Entered STN: 19961022  
Last Updated on STN: 19961022  
Entered Medline: 19961009  
AB The extract from **Sarcocystis** cruzi cysts in bovine muscle was  
subcutaneously injected to mice, guinea pigs, chickens, and rabbits to  
detect its toxicity. Only rabbits showed reactions after administration of  
the extract at a dose of 25 micrograms. The main clinical signs of the  
rabbits were depression, reduction in body temperature and intermittent  
diarrhea and the hematological findings observed were elevation in WBC,  
RBC, PCV, TP, BUN, AST, ALT and creatinine values and reduction in  
glucose, K<sup>+</sup> and pH of blood. The extract, crude toxin, was a water  
soluble, acid-alkali stable and thermolabile protein and estimated to be a  
molecular mass of 15-16 kd.